

A checklist and bibliography of the Opisthobranchia (Mollusca: Gastropoda) of Victoria and the Bass Strait area, south-eastern Australia

ROBERT BURN

Honorary Associate, Museum Victoria, GPO Box 666, Melbourne, Victoria 3001, Australia

Abstract

Burn, R. 2006. A checklist and bibliography of the Opisthobranchia (Mollusca: Gastropoda) of Victoria and the Bass Strait area, south-eastern Australia. *Museum Victoria Science Reports* 10: 1–38.

A checklist of the opisthobranch fauna (Mollusca: Gastropoda: Opisthobranchia) of Victoria and the Bass Strait area, south-eastern Australia comprises a total of 364 nominal species, divided as follows: Acteonida and Cephalaspidea 77 species, Rhodopomorpha 1 species, Sacoglossa 31 species, Anaspidea 10 species, Umbraculida 2 species, Pleurobranchida 6 species, Pteropoda 25 species, and Nudibranchia 212 species. One hundred and thirty eight species (38%) are assigned to genus only; these include both unidentified and unnamed species. Forty species (11%) are yet to be taken alive within the checklist area. The bibliography includes references for the original descriptions of all the genera and all the named species, together with all literature pertinent in some way to the opisthobranch fauna of Victoria, the Bass Strait area, and south-eastern Australia.

Keywords

Mollusca, Gastropoda, Opisthobranchia, Australia, Victoria, Bass Strait, checklist

Introduction

This checklist is a compilation of the 364 opisthobranch mollusc species recorded from, or otherwise known to the author as occurring in, Victoria and the Bass Strait area, south-eastern Australia. Many entries are of unidentified and unnamed species, even generic placement is in doubt in some instances. Geographically, the checklist covers the opisthobranch fauna of the area enclosed by a line in the west from the Victoria/South Australia border to the north-west corner of Tasmania, a line in the east from the Victoria/New South Wales border to the north-east corner of Tasmania, the entire coastline of Victoria along the north, and the north coastline of Tasmania along the south (roughly 37°30–41°S, 140°–150°E). Bass Strait itself is generally less than 100 m deep. However, the area delineated above includes slope waters up to 600 m deep in the west and 2000 m in the east, hence inclusion of some few deep-water species in the checklist.

Zoogeographically, Victoria and the Bass Strait area lies within the southern Australian temperate region, and many opisthobranch species have wide distributions across southern Australia. At times, warmer-water incursions, more commonly from the east than from the west, bring warm-temperate species into the Bass Strait area, and on rarer occasions, tropical and sub-tropical species to eastern Bass Strait. Cool-temperate species from south-eastern Tasmania are also sometimes recorded from Victoria and the Bass Strait area. Further afield, New Zealand too influences the Victorian and Bass Strait fauna with about 20 species in common, not including the pteropods.

Opisthobranchs are found from the intertidal to the deepest seas. The species in this checklist have a depth range of 0–1760 m, with the vast majority occurring in the top 30 m. Because of their beauty, opisthobranchs are favoured animals, especially of the SCUBA-diving fraternity, and many regional guides are available to help identify species seen, photographed or

collected. For the Australian area, at least six such guides have been published within the past 20 years (Willan & Coleman, 1984; Coleman, 1989, 2001; Burn, 1989; Wells & Bryce, 1993; Marshall & Willan, 1999), each of which illustrates some species occurring within the Victoria and Bass Strait area. Most species however are small to very small, often not easily seen in life and difficult to distinguish clearly from their food host, and problematic to photograph and draw. No guide is available to help identify these small species.

Opisthobranchs are a very common component of the intertidal shorelines of Victoria and the Bass Strait area, both in numbers of species and number of specimens. The majority of species are small to very small and are therefore easily overlooked or not recognised. Nevertheless, by exploring the various habitats on and around a rocky reef on the central Victorian coastline (Wilsons Promontory to Cape Otway), it is possible for an experienced person to observe 20–30, exceptionally more than 40, species on a single low tide. Repeated fortnightly observation and exploitation of all accessible habitats over two years at a single location (Point Danger, Torquay, 1980–1981, R. Burn unpublished observations) resulted in a list of 95 species, at least 10% of which were new to science or previously unrecorded for Victoria. Over a longer period and by more than one observer, the channel banks at San Remo, Westernport produced a list exceeding 120 species, again with both new and unrecorded species (Burn, 1990). In deeper waters, the Bass Strait Survey 1979–1984 showed that opisthobranchs are common and diverse throughout the Strait, some 140 live-taken species having been separated from benthic samples. Five stations, sampled by epibenthic sled in depths of 55–130 m, averaged 21 species per station (Burn in Poore et al., 1985). However, large sampling gaps remain to be investigated: little is known of the shallow water opisthobranch fauna of eastern Victoria (Wilsons Promontory to Cape Howe) or of the eastern and western sections of the north coast of Tasmania.

Some species in the checklist have yet to be found alive in Victoria and the Bass Strait area. These species are marked with an asterisk (*). This applies particularly to records of shell-bearing species (Cephalaspidea, Anaspidea, Thecosomata). No attempt has been made to assess the interstitial opisthobranchs that may occur in the marine meiobenthos of south-eastern Australia. Species likely to occur include members of the small order Acochlidioidea, as well as cephalaspideans (*Philine*, *Philinoglossa*) and aeolidioideans (*Pseudovermis*).

The systematic classification followed in the checklist is essentially that tabled in *Mollusca: The Southern Synthesis* (Beesley et al., 1998). Papers published since 1995 (date of the final acceptance of manuscripts for the above publication) and the closing date (30 September 2006) of the checklist present many changes, some major, some minor, some well supported, some speculative. At the time of writing, the newest “Working Classification of Gastropoda” (Bouchet & Rocroi, 2005) summarizes recent phylogenetic research, and speculates upon things to come. Reference to many of these changes is inserted at various points of the checklist.

History of discovery

“The history of discovery of opisthobranchs in Australia is best considered separately for those with shells and those without. Species with shells were included with other shelled molluscs by early collectors and so their discovery follows the path described elsewhere for other marine shells.” (Rudman, 1998: 919). Shell-less species were too much of a problem for early collectors, so much so that they were almost totally ignored. The history of discovery of opisthobranchs in Victoria and the Bass Strait area mirrors that for Australia.

The very first shell-less opisthobranch reported from Victoria and the Bass Strait area was *Pleurobranchaea maculata* by Quoy & Gaimard (1832) who had dredged specimens in Westernport, Victoria between 12th-19th November 1826. Sixty-three years elapsed before the report of another species, *Scyllaea pelagica*, from dredgings near Port Phillip Heads by John Bracebridge Wilson (Hedley, 1895). Then, 10 years later, the great Danish opisthobranch worker Rudolph Bergh described seven northcoast Tasmanian species sent to him by Miss Mary Lodder of Launceston: *Aeolidiella faustina* (= *Spurilla macleayi*), *Alloiodoris marmorata*, *Discodoris dubia* and *D. egea* (= *Paradoris dubia*), *Chromodoris tasmaniensis*, *Aphelodoris luctuosa* (= *A. berghi*), and *Acanthodoris metulifera* (Bergh, 1904; 1905). A very long gap followed.

During the 1930s and 1940s, Mrs Euphemia Freame of Seaholme, well known for her activities in the Field Naturalists Club of Victoria and her small bayside museum, collected opisthobranchs which she forwarded to Joyce Allan at the Australian Museum, Sydney for identification. These identifications were made available to the compilers of, and were included in, an inventory of the marine and estuarine molluscs of Victoria (Macpherson & Chapple, 1951). The

species included in the inventory were: *Ceratosoma brevicaudatum*, *Hoplodoris nodulosa* (as *Staurodoris pustulata*), *Dendrodoris nigra* (as *D. melaena*) *Doriopsilla carneola* (as *Dendrodoris carneola*), *Paradoris dubia* (as *Alloiodoris marmorata*), *Aphelodoris berghi* (as *Archidoris varia*), and *Armina* sp. 1 (as *Armina cygnea* Bergh, 1876, a large species known from NSW, SA and WA, and possibly the northern coastline of Australia). At this stage, the checklist of Victorian and Bass Strait numbered 44 shell-bearing and eight shell-less or nudibranch species, totalling 52 species.

Shortly after, the writer collected his first nudibranch, *Ceratosoma brevicaudatum*, under stones among seagrass, San Remo, Westernport, 13 March 1954. A burgeoning interest soon led to a single-minded concentration on nudibranchs and shelled opisthobranchs, and the rapid expansion of knowledge of the Victorian fauna. The next listing of Victorian marine molluscs (Macpherson & Gabriel, 1962) included 58 shelled and 62 nudibranch species, a total of 120 species. In a little more than 10 years, the known opisthobranch fauna had more than doubled (Burn, 1957a, 1957b, 1958, 1960b). By 1980, the opisthobranch fauna known to the writer from Victoria and the Bass Strait area had more than doubled again to 250 species. This number included many species described or recorded in papers published from 1963 to 1979 (Burn, see bibliography), as well as many species awaiting description for want of additional material and time to work up the descriptive text.

The last quarter century have seen the number increase further with the descriptions of new species (Rudman, 1982, 1983, 1986, 1987a, 1987b, 1990; Willan, 1988; Miller & Willan, 1986). Intensive intertidal and subtidal field work by the writer and colleagues, and deeper water survey work in Bass Strait by Museum Victoria, have revealed additional new records and new species. All species, named, unnamed, and unidentified, have an entry in the checklist, the total now standing at 363, of which 152 are shell-bearing and 212 nudibranch species. And still more species await discovery!

Use of the checklist

Each entry for genus and species in the checklist includes page reference to its original description. In addition, for each species is listed: the type locality where designated; distribution by State and Territory clockwise around Australia (V – Victoria, T – Tasmania, SA – South Australia, WA – Western Australia, NT – Northern Territory, Q – Queensland, NSW – New South Wales; plus NZ – New Zealand, main and subantarctic islands); and depth range. Generic misplacements and species synonyms, by which species have been described or reported in the literature subsequent to the original description, are listed by name only. Reference to recent, mostly colour, figures of species, where available, are also included. In places, some systematic, taxonomic or distributional comment completes the entry. Entries for unnamed or unidentified species include some descriptive comment to separate that species from named congeners, and to aid its recognition.

Checklist**Class Opisthobranchia****Order Acteonida****Superfamily Acteonoidea****Family Acteonidae d'Orbigny, 1843*****Acteon* Montfort, 1810***Acteon* Montfort, 1810: 314*Type species.* *Bulla tornatilis* Linnaeus, 1758****Acteon fructuosus* Iredale, 1936***Acteon fructuosus* Iredale, 1936: 330, pl. xxxiv, fig.28*Type locality.* off Green Cape, NSW, 150 m*Distribution.* NSW, V: 100-150 m****Acteon retusus* Verco, 1907***Acteon retusus* Verco, 1907: 309*Type locality.* Off Beachport, South Australia, 425 m*Distribution.* V, T, SA: 150-200 m*Acteon fructuosus* and *A. retusus* are extremely close in shell characters, and may be forms of the one species.****Acteon subroseus* Iredale, 1936***Acteon subroseus* Iredale, 1936: 330, pl. xxxiv, fig.25*Type locality.* Off Montague Island, NSW, 130-150 m*Distribution.* NSW, V: 100-150 m

Knowledge of the radular characteristics of these three species may lead to their reassignment to other genera within the family.

Obrussena* Iredale, 1930Obrussena* Iredale, 1930: 175(= *Obrussa* Iredale, 1925: 269, preocc.)*Type species.* *Obrussa bracteata* Iredale, 1925

A doubtful inclusion in the family Acteonidae.

****Obrussena bracteata* (Iredale, 1925)***Obrussa bracteata* Iredale, 1925: 269, pl. 42, fig. 15*Type locality.* 140-150 m off Narrabeen*Distribution.* NSW, T: 140-180 m***Pupa* Röding, 1798***Pupa* Röding, 1798: 110*Type species.* *Pupa griselba* Röding, 1798 = *Bulla solidula* Linnaeus, 1758****Pupa affinis* (A. Adams, 1855)***Solidula affinis* A. Adams, 1855: 61*Buccinulus niveus* Angas, 1871a: 19, pl. 1, fig. 27*Type locality.* Moreton Bay, Queensland*Distribution.* Q, NSW, V, SA, WA, tropical and warm temperate Indo-Pacific: 0-50 m

Beu (2004) presents a very extensive synonymy for this presumptive wide ranging tropical and warm temperate species.

Of the two species hitherto listed for Victorian waters, his evidence for the inclusion of *nivea* as a synonym is conclusive. On the contrary, slight differences in the radula support retention of *tragulata* as a separate deeper-water species.****Pupa tragulata* Iredale, 1936***Pupa tragulata* Iredale, 1936: 331, pl. xxiv, fig. 23*Type locality.* Off Sydney, NSW, 160-180 m*Distribution.* NSW, V: 150-200 m**Family Aplustridae Gray, 1847****(=Hydatinidae Pilsbry, 1895)**The well-known family name Hydatinidae Pilsbry, 1895 is invalid. It is a homonym of the earlier "Hydatinidae Ehrenberg, 1838, based on *Hydatina* Ehrenberg, 1828 (Rotatifer); Hydatinidae Ehrenberg is invalid because its type genus is a junior homonym but it remains an available name" (Bouchet & Rocroi, 2005). It is replaced by Aplustridae Gray, 1847, the type genus of which *Aplustrum* Schumacher, 1817 is currently assigned to the synonymy of *Hydatina* Schumacher, 1817 (Rudman, 1972).***Hydatina* Schumacher, 1817***Hydatina* Schumacher, 1817: 186*Type species.* *Bulla physis* Linnaeus, 1758***Hydatina physis* Linnaeus, 1758***Hydatina physis* Linnaeus, 1758: 727*Hydatina physis*.— Wells & Bryce, 1993: 22 (photo: species 5)*Type locality.* unknown*Distribution.* Q, NSW, V, WA, NT, circum-global tropical and warm temperate: 0-10 m**Family Bullinidae Gray, 1850*****Bullina* Férussac, 1822***Bullina* Férussac, 1822: xxx*Type species.* *Bulla scabra* Gmelin, 1791 (non Müller, 1776) = *Bulla lineata* Gray, 1825***Bullina lineata* (Gray, 1825)***Bulla lineata* (Gray, 1825): 408*Bullina lineata*.—Edgar, 1997: 270 (photo)*Type locality.* Sydney, Australia*Distribution.* Q, NSW, V, T, WA, NT, Indo-Pacific tropical and warm temperate: 0-45 m.**Family Ringiculidae Philippi, 1853*****Ringicula* Deshayes, 1838***Ringicula* Deshayes, 1838: 342*Type species.* *Auricula ringens* Lamarck, 1804****Ringicula australis* Hinds, 1844***Ringicula australis* Hinds, 1844: 97*Type locality.* Port Lincoln, South Australia*Distribution.* V: 50-100 m

****Ringicula grandinosa* Hinds, 1844**

Ringicula grandinosa Hinds, 1844: 96

Type locality. Bais Negros, Philippines, 13 m

Distribution. V: 50-100 m

Both *Ringicula australis* and *R. grandinosa* are retainers from earlier lists of Victorian molluscs (eg Macpherson & Gabriel, 1962). Preserved subadult material from deep water in eastern Bass Strait is available but has yet to be formally identified.

Order Cephalaspidea

Superfamily Philinoidea

Family Cylichnidae H. & A. Adams, 1854

***Adamnestia* Iredale, 1936**

Adamnestia Iredale, 1936: 333

Type species. *Adamnestia peroniana* Iredale, 1936 = *Bulla regularis* Gould, 1859 = *Bulla arachis* Quoy & Gaimard, 1833

***Adamnestia arachis* (Quoy & Gaimard, 1833)**

Bulla arachis Quoy & Gaimard, 1833: 361

Bulla regularis Gould, 1859: 140

Cylichna arachis.—Angas, 1867: 226

Adamnestia peroniana Iredale, 1936: 333

Type locality. King George Sound, Western Australia

Distribution. Q, NSW, V, T, SA, WA: 0-200 m

***Cylichna* Lovén, 1846**

Cylichna Lovén, 1846: 10

Type species. *Bulla cylindracea* Pennant, 1777

***Cylichna thetidis* Hedley, 1903**

Cylichna thetidis Hedley, 1903: 395

Type locality. Off Manning River, NSW, 48 m

Distribution. NSW, V, T, SA, WA, NZ: 20-200 m

Gabriel (1962: 206, fig. 4) figured a row of teeth from the radula of *Cylichna thetidis* but inadvertently labelled the figure as representing his new marginellid species *Triginella malinoides* described earlier in the same paper.

***Cylichnella* Gabb, 1873**

Cylichnella Gabb, 1873: 273

Type species. *Bulla bidentata* d'Orbigny, 1841

***Cylichnella* sp**

Distribution. V: 150 m

Ovate shell, columella with strong plait; assignment doubtful.

***Scaphander* Montfort, 1810**

Scaphander Montfort, 1810: 334

Type species. *Bulla lignaria* Linnaeus, 1758

****Scaphander illecebrosus* Iredale, 1925**

Scaphander illecebrosus Iredale, 1925: 269, pl. 42, fig. 14

Type locality. 32 km east of Babel Island, Flinders Island, 120 m

Distribution. T: 150 m

****Scaphander* sp**

Distribution. V: 150 m

A deep water species from off Lakes Entrance, smaller and more trapezoidal in shell shape than *S. illecebrosus* Iredale, 1925.

***Sphaerocylichna* Thiele, 1925**

Sphaerocylichna Thiele, 1925: 242

Type species. *Cylichna atyoides* Thiele, 1925

Sphaerocylichna encompasses a rather uniform series of deep water species, but little is known of the animals.

****Sphaerocylichna incommoda* (E. A. Smith, 1891)**

Bulla incommoda E. A. Smith, 1891: 442

Austrocylichna lagenae Burn, 1978: 99, fig. 9

Type locality. Off Sydney, NSW, 870 m

Distribution. NSW, V, SA: 250-870 m

A probable additional synonym is *Cylichna bulloides* Dell, 1956 from New Zealand.

***Tornatina* A. Adams, 1850**

Tornatina A. Adams in Sowerby, 1850: 554

Types species: *Bulla voluta* Quoy & Gaimard, 1833 (non Gmelin, 1791) = *Tornatina decorata* Pilsbry, 1904

***Tornatina apicina* Gould, 1859**

Tornatina apicina Gould, 1859: 139

Type locality. Sydney Harbour, NSW

Distribution. NSW, V, T: 0-150 m

****Tornatina apiculata* (Tate, 1879)**

Utriculus apiculatus Tate, 1879: 138

Retusa apiculata.—Cotton & Godfrey, 1933: 75

Type locality. King George Sound, Western Australia

Distribution. V, SA, WA: 0-5 m

This species, with type locality King George Sound, WA, is a doubtful record for Victoria.

****Tornatina eumicra* (Crosse in Crosse & Fischer, 1865)**

Bulla eumicra Crosse in Crosse & Fischer, 1865: 40, pl. 12, fig. 7

Retusa eumicra.—Cotton & Godfrey, 1933: 75, pl. 1, fig. 3

Type locality. Spencer Gulf, South Australia

Distribution. V, SA, WA: 0-50 m

This species, with type locality St Vincent Gulf, SA, is a doubtful record for Victoria.

***Tornatina exserta* Hedley, 1903**

Tornatina exserta Hedley, 1903: 393

Type locality. Off Manning River, NSW, 48 m

Distribution. NSW, V, T: 0-200 m

Tornatina sp 1

Distribution. V, SA: 0-20 m

Species distinguished by milky-white blotches in the shell. Records of *Acteocina hofmani* (Angas, 1877) from Victoria are most likely misidentifications of this species. True *hofmani* is larger, more cylindrical, with stronger shoulders.

Tornatina sp 2

Distribution. V: 0-5 m

Exceedingly common estuarine and protected mud-flat species, shell marked by a thick orange-yellow periostracum. This is probably the species identified by Macpherson & Gabriel (1962: 245, fig. 284) as *Acteocina fusiformis* (A. Adams, 1850), a northern Pacific species from Japan.

Tornatina sp 3

Distribution. V, T: 1730 m

Distinguished by very rounded shoulders to the whorls and short nipple-like spire.

Tornatina sp 4

Distribution. V, T: 1730 m

Similar to above species, but with very open aperture.

Family Retusidae Thiele, 1925**Retusa Brown, 1827**

Retusa Brown, 1827: pl. xxxviii, fig. 1

Type species. *Bulla obtusa* Montagu, 1803

***Retusa amphizosta (Watson, 1886)**

Utriculus amphizostus Watson, 1886: 336

Type locality. Cape York, Queensland, 13-17 m

Distribution. Q, NSW, V, T, SA: 0-100 m

May be a complex of similarly shaped species.

Retusa atkinsoni (Tenison Woods, 1876)

Cylichna atkinsoni Tenison Woods, 1876: 156

Type locality. Long Bay, Tasmania

Distribution. NSW, V, T, SA: 0-100 m

Holotype figured by May (1903: 113, fig. 11).

Retusa chrysoma Burn in Burn & Bell, 1974

Retusa chrysoma Burn in Burn & Bell, 1974a: 115-116, figs. 1-3

Type locality. Rocky Point, Yanakie, Corner Inlet, Victoria

Distribution. V: 0-5 m

Retusa atkinsoni and *R. chrysoma* may be forms of the one species.

Retusa pelyx Burn in Burn & Bell, 1974

Retusa pelyx Burn in Burn & Bell, 1974b: 37-38, figs. 1-6

Type locality. Swan Bay, Port Phillip, Victoria

Distribution. V, T: 0-100 m

Retusa protumida (Hedley, 1903)

Cylichna protumida Hedley, 1903: 396

Type locality. Off Cape Three Points, NSW, 85-110 m

Distribution. NSW, V, T, SA: 30-200 m

Examination of the animal of this sharply conical shelled species may prove it to be better placed in *Pyrrunculus* Pilsbry, 1895.

***Retusa pygmaea (A. Adams, 1850)**

Bulla (Cylichna) pygmaea A. Adams, 1850: 595

Type locality. Port Lincoln, South Australia

Distribution. V, T, SA, WA: 0-100 m

Retusa sculpta (Gatliff & Gabriel, 1913)

Bullinella pygmaea sculpta Gatliff & Gabriel, 1913a: 69

Type locality. Off Wilsons Promontory, Victoria

Distribution. V, T: 0-50 m

Retusa sculpta remains enigmatic both for genus and for species. Very few live-taken specimens are known. It would seem that the name *sculpta* applies to the southern temperate-water form of a strongly sculptured tropical species that occurs sparsely along the eastern Australian seaboard.

Retusa sp

Distribution. V: 0 m

A species with regularly ovoid shell showing milky-white patches, the animal selectively feeding upon a different series of foraminiferan to *R. chrysoma* and *R. pelyx* (Burn & Bell, 1974a, b).

Volvulella Newton, 1891

Volvulella Newton, 1891: 268

Type species. *Bulla acuminata* Bruguière, 1792

This is *Rhizorus* auctt. (non Montfort, 1810 = *Bulla* Linnaeus, 1758).

Volvulella rostrata (A. Adams, 1850)

Bulla rostrata A. Adams in Sowerby, 1850: 596

Volvulella parata Iredale, 1936: 332

Type locality. Port Lincoln, South Australia

Distribution. NSW, V, T, SA, WA: 0-200 m

Family Philinidae Gray, 1850**Philine Ascanius, 1772**

Philine Ascanius, 1772: 329

Type species. *Bulla aperta* Linnaeus, 1767

Philine angasi (Crosse in Crosse & Fischer, 1865)

Bullaea angasi Crosse in Crosse & Fischer, 1865: 38

Philine angasi.—Burn, 1989: pl.44.2 (photo)

Philine angasi.—Edgar, 1997: 270, 271 (photo)

Type locality. Spencer Gulf, South Australia

Distribution. Q, NSW, V, T, SA, WA, NZ: 0-500 m

***Philine auriformis* Suter, 1909**

Philine auriformis Suter, 1909: 257

Type locality. Akaroa Harbour, New Zealand, 8-12 m

Distribution. V, T, NZ: 25-90 m

Burn (1969:75) referred to a *Philine* with black-banded gastral plates, collected in 90 m off Lakes Entrance. This has since been identified with the New Zealand *P. auriformis* Suter, 1909, the distribution of which now includes bay localities along the west coast of North America (Gosliner, 1995; Behrens, 2004).

***Philine beachportensis* Verco, 1909**

Philine beachportensis Verco, 1909: 275

Type locality. Off Beachport, South Australia, 365 m

Distribution. V, T, SA: 52-365 m

***Philine columnaria* Hedley & May, 1908**

Philine columnaria Hedley & May, 1908: 123

Type locality. Off Cape Pillar, Tasmania, 220 m

Distribution. NSW, V, T, SA: 0 -200 m

***Philine teres* Hedley, 1903**

Philine teres Hedley, 1903: 398

Type locality. Off Cape Three Points, NSW, 55-110 m

Distribution. NSW, V: 55-150 m

***Philine trapezia* Hedley, 1902**

Philine trapezia Hedley, 1902a: 704

Type locality. Off Shark Point, Sydney Harbour, NSW, 25 m

Distribution. NSW, V: 0-200 m

***Philine* sp 1**

Distribution. V, WA: 0-50 m

Orange-bodied, epifaunal species, possibly related to the widespread Indo-Pacific *P. rubrata* Gosliner, 1988.

***Philine* sp 2**

Distribution. V: 0 m

Minute (2-3 mm long), white body, epifaunal.

***Philine* sp 3**

Distribution. V, T, SA: 0 m

White cylindrical body, anterior shell margin with long denticles; infaunal.

***Philine* sp 4**

Distribution. NSW, V: 0-3 m

Small, anteriorly slender, elongate animal, head much longer than visceral hump, infaunal in estuaries.

Family Aglajidae Pilsbry, 1895

***Melanochlamys* Cheeseman, 1881**

Melanochlamys Cheeseman, 1881: 224

Type species. *Melanochlamys cylindrica* Cheeseman, 1881

***Melanochlamys queritor* (Burn, 1957)**

Aglaja queritor Burn, 1957a:115

Aglaja henri Burn 1969: 71, figs. 8-9

Aglaja (Melanochlamys) queritor.— Burn, 1974: 50

Type locality. Portarlington, Port Phillip, Victoria

Distribution. NSW, V, SA: 0-6 m

***Melanochlamys* sp**

Melanochlamys sp. — Coleman, 2001: 119 (photo)

Distribution. V, T, SA: 3-71 m

Pale greyish body with sparse brown spotting dorsally; figured (Coleman, 2001: 119) as *Melanochlamys* sp from Bass Strait, 3 m on sand.

***Noalda* Iredale, 1936**

Noalda Iredale, 1936: 334

Type species. *Hydatina exigua* Hedley, 1912

***Noalda exigua* (Hedley, 1912)**

Hydatina exigua Hedley, 1912: 158

Noalda exigua. — Burn, 1998: 953, fig. 16-31 C-F

Type locality. Middle Head, Sydney Harbour, NSW

Distribution. NSW, V, T: 0-37 m

Animal and shell figured (Burn, 1998) from Point Danger, Torquay, Victoria, on algae at low tide.

***Philinopsis* Pease, 1860**

Philinopsis Pease, 1860a :21

Type species. *Philinopsis speciosa* Pease, 1860

***Philinopsis cyanea* (Martens, 1879)**

Doridium cyanea Martens, 1879: 738

Type locality. Inhambane, Mozambique

Distribution. Q, NSW, V, WA, NT, tropical to warm temperate Indo-west Pacific: 0-20 m

Rudman (2006; “March 24. Comment on *Philinopsis cyanea*? from Victoria”) identifies, and pictures a 25 mm long animal seen at Steeles Rock, Portarlington, Port Phillip, Victoria by Trevor McMurrich, 18 March 2006. Reference to the original notes and sketches of the specimen reported (Burn, 1957a: 117) as *Aglaja taronga* from Swan Bay, Port Phillip, Victoria strongly suggests that this specimen should be re-identified as *Philinopsis cyanea*. The 1957 specimen was velvet black in colour with a narrow white edging to the parapodia, an orange line each side of the anterior part of the head shield, an orange submarginal band along each parapodium, and the posterior end of the head shield was held abruptly raised. This matches almost exactly the specimen illustrated in Rudman (2006).

***Philinopsis lineolata* (H. & A. Adams, 1854)**

Aglaja lineolata H.& A. Adams, 1854: 27, pl. 58, fig. 4

Philinopsis lineolata.—Wells & Bryce, 1993: 31, 33 (photo: species 22)

Philinopsis lineolata.—Coleman, 2001: 120 (photo)

Type locality. Indo-Pacific

Distribution. Q, NSW, V, SA, WA, NT, NZ, tropical and temperate Indo-Pacific: 0-60 m

***Philinopsis taronga* (Allan, 1933)**

Aglaiia taronga Allan, 1933: 444
Aglaja (Philinopsis) taronga.— Burn, 1974: 50
Chelidonura aureopunctata Rudman, 1968: 221
Philinopsis taronga.— Coleman, 2001:121 (photo)

Type locality. Athol Bay, Sydney Harbour, NSW

Distribution. NSW, V, T, NZ: 0-150 m

***Philinopsis* sp 1**

Distribution. V: 0 m

Distinguished from congeners by the presence of short tentaculiform corners of the anterior foot.

***Philinopsis* sp 2**

Distribution. V: 25-35 m

A small (<5 mm) all white species from deeper water off the Gippsland coast. Possibly to be identified with *Philinopsis virgo* (Rudman, 1968) from 100 m in northern New Zealand, a larger (20 mm) completely white species.

***Philinopsis* sp 3**

Distribution. V, SA, WA: 5-40 m

Pale brown species with network of brown lines on foot.

Family Gastropteridae Swainson, 1840***Gastropteron* Kosse, 1813**

Gastropteron Kosse, 1813: 10

Type species. *Gastropteron meckeli* Kosse, 1813

***Gastropteron* sp**

Distribution. V, T: 70-150 m

Animal with large, very thin external shell enclosing visceral mass.

***Sagaminopteron* Tokioka & Baba, 1964**

Sagaminopteron Tokioka & Baba, 1964: 218

Type species. *Sagaminopteron ornatum* Tokioka & Baba, 1964

***Sagaminopteron ornatum* Tokioka & Baba, 1964**

Sagaminopteron ornatum Tokioka & Baba, 1964: 218
Sagaminopteron ornatum.—Wells & Bryce, 1993: 29 (photo: species 17)
Sagaminopteron ornatum.—Edgar, 1997: 271 (photo)
Sagaminopteron ornatum.—Coleman, 2001: 121 (photo)

Type locality. Sagami Bay, Japan, 13 m

Distribution. Q, NSW, V, SA, WA, tropical and temperate Indo-Pacific: 0-25 m

***Siphopteron* Gosliner, 1989**

Siphopteron Gosliner, 1989: 340

Type species. *Siphopteron tigrinum* Gosliner, 1989

***Siphopteron* sp 1**

Distribution. V, T: 0-20 m

An all orange or orange and yellow species with black tip to cephalic siphon and to visceral appendage.

***Siphopteron* sp 2**

Distribution. V, WA: 0-20 m

Red or orange body with longitudinal blue lines on outer surfaces; close to *S. tigrinum* Gosliner, 1989.

***Siphopteron* sp 3**

Distribution. V: 0 m

A dark greyish species with pattern of reddish tessellated patches on visceral mass and parapodia.

Superfamily Bulloidea**Family Bullidae Gray, 1827*****Bulla* Linnaeus, 1758**

Bulla Linnaeus, 1758: 725

Type species. *Bulla ampulla* Linnaeus, 1758

***Bulla quoyii* Gray in Dieffenbach, 1843**

Bulla australis Gray, 1825: 408 (non Férussac, 1822)
Bulla australis Quoy & Gaimard, 1833: 357 (non Férussac, 1822)
Bulla quoyii Gray in Dieffenbach 1843: 243
Bulla tenuissima Sowerby, 1868: pl. 1, fig. 4
Bullaria botanica Hedley, 1918: M104
Quibulla botanica.—Iredale, 1929: 349
Bulla quoyii.—Willan, 1978: 58
Bulla quoyii.—Wells & Bryce, 1993: 25 (photo: species 10)
Bulla quoyii.—Edgar, 1997: 283 (photo)

Type locality. New Zealand

Distribution. Q, NSW, V, T, SA, WA, NZ: 0-20 m

Willan (1978) reviewed the nomenclatural complexities of this species, and introduced the name *Bulla quoyii* into the Australian molluscan fauna. Wells (1985) added *Bulla tenuissima* Sowerby, 1868 to the synonymy of *Bulla quoyii*.

Family Haminoeidae Pilsbry, 1895***Austrocylichna* Burn, 1974**

Austrocylichna Burn, 1974a: 44

Type species. *Bulla exigua* A. Adams, 1850

***Austrocylichna exigua* (A. Adams, 1850)**

Bulla exigua A. Adams, 1850 in Adams, 1850: 589

Type locality. Port Lincoln, South Australia

Distribution. V, T, SA, WA: 0-20 m

***Cylichnatys* Kuroda & Habe, 1952**

Cylichnatys Kuroda & Habe, 1952: 51

Type species. *Bullinella striata* Yamakawa, 1911 = *Haminea angusta* Gould, 1859

***Cylichnatys campanula* Burn, 1978**

Cylichnatys campanula Burn, 1978b: 104-106, figs. 11-17

Type locality. Rocky Point, Yanakie, Corner Inlet, Victoria

Distribution. NSW, V, T, SA, WA: 0-5 m

***Haminoea* [Turton] in Turton & Kingston in Carrington, 1830**

Haminoea [Turton] in Turton & Kingston in Carrington 1830: genus no. 63 (signature F₈)

Type species. *Bulla hydatis* Linnaeus, 1758

ICZN Opinion 1942 (2000) corrects the spelling of both family and genus names to that used above, and attributed the genus name to an originally anonymous contribution by Turton, indicated by the square brackets, [], enclosing that author's name.

***Haminoea maugeansis* Burn, 1966**

Haminoea maugeansis Burn, 1966c: 330-331, figs 1-2
Haminea tenera.—Angas, 1871b: 98 (*non* A. Adams, 1850)

Type locality. Port MacDonnell, South Australia

Distribution. V, T, SA: 0-22 m

***Haminoea* sp**

Distribution. V, SA, WA: 0 m

A small, thin shelled species, animal cream or yellowish without dark pigmented rosettes in the shell mantle.

***Liloa* Pilsbry, 1921**

Liloa Pilsbry, 1921: 370

Type species. *Haminea tomaculum* Pilsbry, 1917

***Liloa brevis* (Quoy & Gaimard, 1833)**

Bulla brevis Quoy & Gaimard, 1833: 358
Haminea brevis.—Angas, 1865: 188
Bulla cuticulifera Smith, 1872: 350
Liloa brevis.—Burn, 1989: pl.43.6 (photo)
Liloa brevis.—Coleman, 2001: 123 (photo)

Type locality. King George Sound, Western Australia

Distribution. Q, NSW, V, T, SA, WA: 0-22 m

***Limulatys* Iredale, 1936**

Limulatys Iredale, 1936: 328

Type species. *Limulatys reliquus* Iredale, 1936

***Limulatys reliquus* Iredale, 1936**

Limulatys reliquus Iredale, 1936: 328, pl.24, fig. 20

Type locality. Sydney Harbour, NSW, dredged

Distribution. NSW, V, NZ: 0-35 m

***Nipponatys* Kuroda & Habe, 1952**

Nipponatys Kuroda & Habe, 1952: 72

Type species. *Alicula volvulina* A. Adams, 1862

****Nipponatys tumidus* Burn, 1978**

Nipponatys tumidus Burn, 1978b: 101-102, fig. 10

Type locality. Thompsons Creek, Breamlea, Victoria

Distribution. V, T: 0-20 m

Live-taken specimens are known only from south-eastern Tasmania

Superfamily Diaphanoidea**Family Diaphanidae Odhner, 1914*****Colpodaspis* M. Sars, 1870**

Colpodaspis Sars, 1870: 70-74

Type species. *Colpodaspis pusilla* M. Sars, 1870

***Colpodaspis* sp 1**

Distribution. V: 0-115 m

Dark blue mantle, without siphonal fold.

***Colpodaspis* sp 2**

Distribution. V, SA: 0-95 m

Dark brown mantle, with siphonal fold.

***Colpodaspis* sp 3**

Colpodaspis sp 3 Wells & Bryce, 1993: 24 (photo: species 9), 25

Distribution. V, WA: 0-15 m

Pale blue mantle, with siphonal fold.

***Diaphana* Brown, 1827**

Diaphana Brown, 1827: pl. 38

Type species. *Diaphana candida* Brown, 1827 = *Diaphana minuta* Brown, 1827

***Diaphana brazieri* Angas, 1877**

Diaphana brazieri Angas, 1877: 175, pl. 26, fig. 20
Austrodiaphana brazieri Pilsbry, 1895: 287, pl. 26, fig. 68
Aplustrum brazieri Hedley, 1902b: 16, pl. 3, fig. 36

Type locality. Sow & Pigs Reef, Sydney Harbour, NSW, 9 m

Distribution. Q, NSW, V, T, SA, WA, NZ, ?Japan: 0-600 m

Schiøtte (1998) synonymizes the three New Zealand species, *Austrodiaphana colei* Fleming, 1948, *A. maunganuica* Powell, 1952 and *A. flemingi* Powell, 1955, and doubtfully the Japanese *Diaphana sakurarii* Habe, 1976, with the Australian *Diaphana brazieri*.

***Diaphana tasmanica* (Beddome, 1883)**

Akera tasmanica Beddome, 1883: 169

Type locality. Off Old Station, Browns River Road, Tasmania, 15 m

Distribution. V, T, NZ: 0-10 m

Shell figured in Gatliff and Gabriel (1908a: pl. 21, fig. 6-7), May (1923a: pl. 46, fig. 15) and Schiøtte (1998: fig. 7D-H, 8B).

***Diaphana* sp**

Distribution. V: 70-200 m

A globose species similar in shell shape to *D. abyssalis* Schiøtte, 1998 and *D. globosa* (Lovén, 1846) (Schiøtte, 1998).

***Rhinodiaphana* Lemche, 1967**

Rhinodiaphana Lemche, 1967: 208

Type species. *Utriculus ventricosus* Jeffreys, 1865

Rhinodiaphana sp

Distribution. V: 130 m

A deep-water species from eastern Bass Strait, shell external with very wide aperture as in the Northern European type species.

Toledonia Dall, 1902

Toledonia Dall, 1902: 512

Type species. *Toledonia perplexa* Dall, 1902

Toledonia sp

Distribution. V: 25 m

Closely related to *T. succineaformis* Powell, 1955 from off the Auckland Islands, south of New Zealand.

Superfamily Runcinoidea**Family Runcinidae H. & A. Adams, 1854****Runcina Forbes in Forbes & Hanley, 1851**

Runcina Forbes in Forbes & Hanley, 1851: 611

Type species. *Runcina hancocki* Forbes in Forbes & Hanley, 1851 = *Pelta coronata* Quatrefages, 1844

Runcina australis Burn, 1963

Runcina australis Burn, 1963a: 11-14, figs 1-11

Type locality. Point Danger, Torquay, Victoria

Distribution. NSW, V, SA: 0-5 m

Runcina sp 1

Distribution. V: 0-51 m

Distinguished by the presence of a minute, partly coiled external shell projecting from the posterior notum.

Runcina sp 2

Distribution. V: 30-70 m

A deeper water species somewhat similar to the New Zealand *Runcinella zelandica* Odhner, 1924.

Family Ilbiidae Burn, 1963**Ilbia Burn, 1963**

Ilbia Burn, 1963a: 15

Type species. *Ilbia ilbi* Burn, 1963

Ilbia ilbi Burn, 1963

Ilbia ilbi Burn, 1963a: 15-18, fig. 29-30
Ilbia ilbi.—Coleman, 2001: 123 (photo)

Type locality. Point Lonsdale, Victoria

Distribution. NSW, V: 0-12 m

Order Rhodopemorpha**Family Rhodopidae Ihering, 1876**

Systematic position unresolved.

Rhodope Koelliker, 1847

Rhodope Koelliker, 1847: 239

Type species. *Rhodope veranii* Koelliker, 1847

Haszprunar & Hess (2005) comparatively review all described and undescribed *Rhodope* species.

Rhodope sp

Distribution. V: 0 m

Known only from a single living specimen (Burn, 1998: 961, fig. 16.43).

Order Sacoglossa**Superfamily Oxynooidea****Family Volvatellidae Pilsbry, 1895****Ascobulla Ev. Marcus, 1972**

Ascobulla Ev. Marcus, 1972: 286

Type species. *Cylindrobulla ulla* Er. Marcus & Ev. Marcus, 1970

Ascobulla fischeri (A. Adams & Angas, 1864)

Cylindrobulla fischeri A. Adams & Angas, 1864: 37

Ascobulla fischeri.—Wells & Bryce, 1993: 59 (photo: species 59)

Type locality. Spencer Gulf, South Australia

Distribution. NSW, V, T, SA, WA: 0-15 m

Family Oxynoidae Stoliczka, 1868**Oxynoe Rafinesque, 1814**

Oxynoe Rafinesque, 1814a: 162

Type species. *Oxynoe olivacea* Rafinesque, 1814

Oxynoe viridis (Pease, 1861)

Lophocercus viridis Pease, 1861: 246

Oxynoe viridis.—Burn, 1989: pl.46.5 (photo)

Oxynoe viridis.—Wells & Bryce, 1993: 61 (photo: species 63), 62

Oxynoe viridis.—Edgar, 1997: 274 (photo)

Oxynoe viridis.—Coleman, 2001:128 (photo)

Type locality. Pacific Islands

Distribution. Q, NSW, V, T, SA, WA, tropical and temperate Indo-Pacific: 0-10 m.

Roburnella Ev. Marcus, 1982

Roburnella Ev. Marcus, 1982:15

Type species. *Lobiger wilsoni* Tate, 1889

Roburnella wilsoni (Tate, 1889)

Lobiger wilsoni Tate, 1889: 66, pl.11, fig.12

Lophopleurella wilsoni Burn, 1966: 58

Roburnella wilsoni.—Burn, 1989: pl.46.4 (photo)

Roburnella wilsoni.—Coleman, 2001:129 (photo)

Type locality. South Channel, Port Phillip, Victoria, 16-34 m

Distribution. V, T, SA, WA: 0-22 m.

Family Juliidae E.A. Smith, 1885

Overseas workers have assigned all thin-shelled bivalved gastropods to the oldest available genus, *Berthelinia* Crosse, 1875, based upon the Paris Basin Eocene fossil *Berthelinia elegans* Crosse, 1875. The writer accepts that *Berthelinia* and *Tamanovalva* are probably synonymous, but retains *Edentellina* and *Midorigai* as separate genera. A fourth species

of this family occurring in Victorian waters appears to belong to yet another genus; provisionally, it is listed as *Berthelinia* sp.

***Edenttellina* Gatliff & Gabriel, 1911**

Edenttellina Gatliff & Gabriel, 1911: 190

Type species. *Edenttellina typica* Gatliff & Gabriel, 1911

***Edenttellina typica* Gatliff & Gabriel, 1911**

Edenttellina typica Gatliff & Gabriel, 1911: 190, pl.46, figs 5-6

Edenttellina typica.— Burn, 1989: pl.46.3 (photo)

Edenttellina typica.— Coleman, 2001: 127 (photo)

Type locality. Portsea, Port Phillip, Victoria

Distribution. V, T, SA: 0-2 m

***Midorigai* Burn, 1960**

Midorigai Burn, 1960b: 45

Type species. *Midorigai australis* Burn, 1960: 46

***Midorigai australis* Burn, 1960**

Midorigai australis Burn, 1960b: 46, figs 8-14

Midorigai australis.— Burn, 1989: pl.46.2 (photo)

Midorigai australis.— Coleman, 2001:127-128 (photo)

Type locality. Torquay, Victoria

Distribution. V, T, SA: 0-2 m

Berthelinia rotnesti (Jensen, 1993), from Rottnest Island, Western Australia, is the western cognate of, if not identical with, *Midorigai australis*. Both species have an obligate association with the green alga, *Caulerpa simpliciuscula*.

***Tamanovalva* Kawaguti & Baba, 1959**

Tamanovalva Kawaguti & Baba, 1959: 178

Type species. *Tamanovalva limax* Kawaguti & Baba, 1959

***Tamanovalva babai* Burn, 1965**

Tamanovalva babai Burn, 1965b: 735

Tamanovalva babai.— Burn, 1989: pl.46.1 (photo)

Type locality. Point Danger, Torquay, Victoria

Distribution. V, T, SA: 0-3 m

***Berthelinia* sp**

Distribution. V, WA: 0-2 m

Shell mantles plain green, shell shorter and broader than *Tamanovalva babai*, protoconch smaller and less upright.

Superfamily Plakobranchoidea

Family Plakobranchoidea Gray, 1840

***Elysia* Risso, 1818**

Elysia Risso, 1818: 375

Type species. *Elysia timida* Risso, 1818

***Elysia australis* (Quoy & Gaimard, 1832)**

Actaeon australis Quoy & Gaimard, 1832: 317

Type locality. Port Jackson, NSW

Distribution. NSW, V: 0-2 m

Thin whip-like rhinophores and a broader body separate *Elysia australis* from its partially sympatric congener *E. coodgensis*,

which has cylindrical rhinophores and an always present black stripe on the head.

***Elysia coodgensis* Angas, 1864**

Elysia coodgensis Angas, 1864: 69, pl. 6, fig 4

Elysia australis.— Basedow & Hedley, 1905: 148 & subsequent authors (*non* Quoy & Gaimard, 1832)

Elysia australis.— Wells & Bryce, 1993: 61 (photo: species 64), 62

Elysia coodgensis.— Coleman, 2001:129 (photo)

Type locality. Coodgee Bay, Sydney, NSW

Distribution. Q, NSW, V, SA, WA, NT: 0-3 m

***Elysia furvacauda* Burn, 1958**

Elysia furvacauda Burn, 1958: 5, pl. 6, fig 1

Type locality. Torquay, Victoria

Distribution. V: 0-3 m

A dark green or reddish-brown species densely speckled with reddish orange and blue pigment cells, with cream pigment marking the sinuses of the parapodial margins, the rhinophores auriculate, and the foot corners angulate.

***Elysia maoria* Powell, 1937**

Elysia maoria Powell, 1937: 121, pl.30, fig. 5

Elysia maoria.— Coleman, 2001:129 (photo)

Type locality. Takapuna Reef, Auckland, New Zealand

Distribution. Q, NSW, V, NZ: 0-5 m

A dark green species associated with *Codium fragile tomentosoides* upon which it lives and feeds.

***Elysia* sp 1**

Distribution. V: 0 m

A reddish brown species with two elongate tongue-like lobes projecting from each parapodial edge.

***Elysia* sp 2**

Distribution. V: 0 m

A small green species with two sharp black-tipped projections along each parapodial edge.

***Elysia* sp 3**

Distribution. V, SA: 0 m

This species was confused by Burn (1990) with *Elysia furvacauda* (Burn, 1990). Additional material of both this species and *E. furvacauda* indicates their separation, and that neither species is *E. japonica*, as suggested by Jensen (1985). This is a green or brownish species with black rhinophoral and tail tips. It is periodically common among seagrass and algae at San Remo, Westernport.

Superfamily Limapontioidea

Family Caliphyllidae Tiberi, 1881

***Polybranchia* Pease, 1860**

Polybranchia Pease, 1860b: 141

Type species. *Polybranchia pellucida* Pease, 1860

***Polybranchia pallens* (Burn, 1957)**

Cyerce nigra pallens Burn, 1957b: 14, pl. 3, figs 8-11
Polybranchia pallens.— Burn, 1989: pl.47.3 (photo)
Polybranchia pallens.— Coleman, 2001:134 (photo)

Type locality. Queenscliff, Victoria

Distribution. V, T, SA: 0-10 m

Family Limapontiidae Gray, 1847***Costasiella Pruvot-Fol, 1951***

Costasiella Pruvot-Fol, 1951: 73

Type species. *Costasiella virescens* Pruvot-Fol, 1951

***Costasiella* sp 1**

Distribution. NSW, V: 0 m

Drab yellowish-brown animal with high-domed pericardium and tentaculiform foot-corners.

***Costasiella* sp 2**

Distribution. V: 0 m

Drab yellowish-brown animal with dorsally flattened pericardial swelling and rounded anterior foot.

***Aplysiopsis* Deshayes, 1853**

Aplysiopsis Deshayes, 1853: explanation of plates p.56

Type species. *Aplysiopsis elegans* Deshayes, 1853

***Aplysiopsis formosa* Pruvot-Fol, 1953**

Aplysiopsis formosa Pruvot-Fol, 1953: 47, pl. ii, fig. 21
Aplysiopsis formosa.— Coleman, 2001: 133 (photo)

Distribution. V, Mediterranean, Florida, temperate North Atlantic: 0-12 m.

Type locality. Temara, Morocco

A dark green species with a pair of black stripes showing lengthways on the sole; lives on and eats the green alga *Cladophora prolifera*.

***Ercolania* Trinchese, 1872**

Ercolania Trinchese, 1872: 86

Type species. *Ercolania siotti* Trinchese, 1872

***Ercolania margaritae* Burn, 1974**

Ercolania margaritae Burn, 1974a: 52, figs 7-10

Type locality. Point Lonsdale, Victoria

Distribution. NSW, V, T: 0-16 m

***Ercolania* sp 1**

Distribution. V: 0 m

A bright green and pink species living on the green alga *Apjohnia laetivirens*.

***Ercolania* sp 2**

Distribution. V: 0 m

A mottled brownish species with very short rhinophores, and anal papilla projecting from the right side of the renal ridge well behind the pericardium. A very juvenile, newly settled

specimen, less than 0.5 mm long, of this species was the basis of the record of *Limapontia* sp. from Victorian (Burn, 1973b).

***Ercolania* sp 3**

Distribution. V: 0 m

A compact very black species with white stripe on each slender rhinophore, white tips to cerata, and anus flat on pericardium.

***Ercolania* sp 4**

Distribution. V: 0-6 m

A black species with long slender white rhinophores, red or reddish-brown tips to cerata, and anus flat on pericardium. This species very closely resembles some figures of the Japanese *Ercolania boodleae* (Baba, 1938) (Nakano, 2004: 56, fig. 113).

***Hermaea* Lovén, 1844**

Hermaea Lovén, 1844: 50

Type species. *Doris bifida* Montagu, 1815

***Hermaea* sp 1**

Hermaea sp.— Coleman, 2001: 133 (photo: 'Pink Hermaea')

Distribution. NSW, V, SA: 0-10 m

Brown line on pharynx, large leaf-life cerata, red digestive gland, high anal papilla.

***Hermaea* sp 2**

Hermaea sp.— Coleman, 2001: 133 (photo: 'Inflated Hermaea')

Distribution. V: 0-10 m

Cerata swollen, knobably, club-shaped, red digestive gland, high anal papilla.

***Hermaea* sp 3**

Distribution. V: 0 m

Cerata slender, knobably, club-shaped, green digestive gland, anus low on neck.

***Hermaea* sp 4**

Distribution. V: 0 m

Cerata swollen, knobably with green digestive gland, anus low on neck.

***Hermaea* sp 5**

Distribution. V: 0 m

Rhinophores simple (not auriculate as in other four species), cerata slender, knobably, with very pale digestive gland.

***Placida* Trinchese, 1876**

Placida Trinchese, 1876: 84

Type species. *Laura viridis* Trinchese, 1873

***Placida dendritica* (Alder & Hancock, 1843)**

Hermaea dendritica Alder & Hancock, 1843: 233

Placida dendritica.— Burn, 1989: pl. 47.4 (photo)

Placida dendritica.— Coleman, 2001: 133 (photo)

Type locality. Torbay, England

Distribution. Q, NSW, V, T, SA, WA: 0-10 m

This circumglobal *Codium*-eating species probably represents a species complex, the members of which have still to be sorted out. The earliest name for an Australasian species is *Stiliger aoteana* Powell, 1937.

***Placida* sp**

Distribution. V: 0-6 m

Differs from local specimens assigned to *Placida dendritica* by larger size, paler colour, different branching of the digestive gland to the cerata, and a different food alga (*Bryopsis*).

***Stiliger* Ehrenberg, 1828**

Stiliger Ehrenberg, 1828: pl. 1, fig. 3

Type species. *Stiliger ornatus* Ehrenberg, 1828

***Stiliger smaragdinus* Baba, 1949**

Stiliger smaragdinus Baba, 1949: 32, 129

Stiliger smaragdinus.— Edgar, 1997: 275 (photo)

Stiliger smaragdinus.— Vafiadis, 1999: 118 (photo)

Stiliger smaragdinus.— Coleman, 2001: 133 (photo)

Type locality. Off Sajima, Sagami Bay, Japan, 16 m

Distribution. Q, NSW, V, T, SA, WA, NT, NZ, tropical and temperate Indo-Pacific: 0-10 m

Order Anaspeidea

Within this order, Willan (1998) and Valdés & Bouchet (2005) used two superfamilies, Akerioidea and Aplysioidea, each with a single family, Akeridae and Aplysiidae. Molecular, morphological and histological re-assessment of genus-level taxa (Medina & Walsh, 2000; Klussman-Kolb, 2004) confirmed the placement, basally, of *Akera* within the order, and indicated hitherto unrecognised relationships at both genus and species level.

Family Akeridae Mazzarelli, 1891

***Akera* Müller, 1776**

Akera Müller, 1776: 242

Type species. *Akera bullata* Müller, 1776

****Akera soluta* (Gmelin, 1791)**

Bulla soluta Gmelin, 1791: 3434

Akera solute Gatliff & Gabriel, 1908b: 386

Akera soluta.— Willan, 1998: 975, fig. 16.56

Akera soluta.— Coleman, 2001: 123 (photo)

Type locality. Zanzibar

Distribution. Q, NSW, V, WA, tropical and temperate Indo-Pacific: 0-10 m

Reported from San Remo, Victoria by Gatliff & Gabriel (1908: 386) from specimens in the collection of Mrs Agnes F. Kenyon of Melbourne, a well-known late 19th century shell-collector whose collection was purchased by Adelaide surgeon and malacologist Sir Joseph Verco, now deposited in the South Australian Museum. Mrs Kenyon did not include *Akera soluta* in her "A list of marine Mollusca of Victoria" (1898: 1-12, privately published: Melbourne). The only specimens (two dry lots) in the Museum Victoria collection are both localized as San Remo, Victoria, and are part of a large molluscan collection donated by the Queen Victoria Museum, Launceston, Tasmania in 1948. These are unaccompanied by collector or collection data. *Akera soluta* may occur at times in the estuaries of the far

east of the State; the nearest live-taken records are from Merimbula Inlet, southern NSW (Day & Hutchings, 1984).

Family Aplysiidae Lamarck, 1809

***Aplysia* Linnaeus, 1767**

Aplysia Linnaeus, 1767: 1072

Type species. *Aplysia depilans* Gmelin, 1791

***Aplysia dactylomela* Rang, 1828**

Aplysia dactylomela Rang, 1828b: 56

Aplysia dactylomela.— Wells & Bryce, 1993: 43, 44 (photo: species 39)

Aplysia dactylomela.— Edgar, 1997: 272, 273 (photo)

Type locality. Cape Verde Islands

Distribution. Circum-global tropical and temperate seas: 0-20 m

***Aplysia juliana* Quoy & Gaimard, 1832**

Aplysia juliana Quoy & Gaimard, 1832: 309

Type locality. Mauritius

Distribution. Circum-global tropical and temperate seas: 0-20 m

The record of *Aplysia nigra* d'Orbigny, 1835 from Portland (Macpherson & Gabriel, 1962: 248) is possibly a misidentification of *Aplysia juliana*.

***Aplysia parvula* Guilding in Mörch, 1863**

Aplysia parvula Guilding in Mörch, 1863: 22

Aplysia concava Sowerby, 1869: pl. 6, species 24

Aplysia norfolkensis Sowerby, 1869: pl. 10, species 42

Aplysia parvula.— Wells & Bryce, 1993: 43, 44 (photo: species 38)

Aplysia parvula.— Edgar, 1997: 272 (photo)

Type locality. St Thomas, West Indies

Distribution. Circum-global tropical and temperate seas: 0-50 m

***Aplysia sowerbyi* Pilsbry, 1895**

Aplysia sowerbyi Pilsbry, 1895: 101

Type locality. Sydney, NSW

Distribution. Q, NSW, V: 0-2 m

***Aplysia sydneyensis* Sowerby, 1869**

Aplysia sydneyensis Sowerby, 1869: pl. 7, fig. 31

Aplysia sydneyensis.— Edgar, 1997: 273 (photo)

Type locality. Sydney, NSW

Distribution. Q, NSW, V, T: 0-23 m

***Bursatella* Blainville, 1817**

Bursatella Blainville, 1817: 138

Type species. *Bursatella leachii* de Blainville, 1817

***Bursatella leachii* Blainville, 1817**

Bursatella leachii Blainville, 1817: 138

Bursatella leachii.— Coleman, 2001: 125 (photo)

Type locality. Indian Ocean

Distribution. Circum-global tropical and temperate seas: 0-30 m

A new record for Victoria. A single specimen of the NSW form of this species, collected in Western Port, is all that is known for Victoria (MV F19109). Coleman (2001:127, Southern Sea Hare) figured a 48 mm long animal from Mallacoota, 3 m on

reef, as *Petalifera* sp. This appears to be a small *Bursatella leachii*.

***Bursatella* sp**

Bursatella leachi.— Wells & Bryce, 1993: 46, 47 (photo: species 44) (*non* Blainville, 1817)
Bursatella sp.— Edgar, 1997: 273 (photo)

Distribution. V, SA, WA: 0-10 m

A new record for Victoria; recently (6 May 2006) observed and photographed alive at Blairgowrie boat harbour, Port Phillip. Separated from the NSW form of *Bursatella leachii* by the dense coat of slender pointed papillae, the long narrow neck, and the humped, rather than swollen, body. Appears to be endemic to south Western Australia and South Australia, with rare incursions eastward into Victorian waters.

***Dolabrifera* Gray, 1847**

Dolabrifera Gray, 1847: 162

Type species. *Dolabella dolabrifera* Cuvier, 1804

***Dolabrifera brazieri* Sowerby, 1870**

Dolabella brazieri Sowerby, 1870: 250

Type locality. Northhead, Botany Bay, NSW

Distribution. NSW, V, NZ: 0-5 m

A new record for Victoria. In March 2005, a large 150 mm long specimen was seen and photographed by John Eichler at Cape Conran, eastern Victoria. In late February 2006, three slightly smaller 125 mm long specimens, two mottled pale fawn with wide green foot margin and one entirely dark brown with narrower yellowish foot margin, together with several typical flat parallel-stitched egg ribbons under stones, were observed and photographed at the same place by John Eichler and Platon Vafiadis. One specimen deposited in the Museum Victoria collection (F110093). Size and proximity to its south-eastern Australian range identifies these specimens as *Dolabrifera brazieri* Sowerby, 1870. Otherwise, there is little to separate *D. brazieri* from the smaller (to 50 mm long) circum-global tropical and temperate species *D. dolabrifera* (Cuvier, 1817); Klussman-Kolb (2004) examined specimens of both species, and maintained them under two names.

***Petalifera/Phyllaplysia* sp.**

Distribution. V: 2 m

A small flat greenish species is sometimes common on seagrass (*Posidonia*) in the inlets of eastern Victoria. It remains to be identified at both genus and species level.

Order Umbraculida

Wägale & Willan (2000) found the Pleurobranchioidea to be the sister-group to the Nudibranchia, proposing the new higher taxon Nudipleura to encompass both groups. Consequently, Notaspidea retained only the superfamily Umbraculoidea. Since then, Notaspidea has been replaced by Umbraculida (Valdés & Bouchet, 2005).

Superfamily Umbraculoidea

Family Tylodinidae Gray, 1847

***Tyrodina* Rafinesque, 1814**

Tyrodina Rafinesque, 1814a: 162

Type species. *Tyrodina punctulata* Rafinesque, 1814 = *Patella perversa* Gmelin, 1791

***Tyrodina corticalis* (Tate, 1889)**

Umbrella corticalis Tate, 1889: 65, pl. 11, fig. 11
Tyrodina corticalis.— Burn, 1989: pl. 47.5 (photo)
Tyrodina corticalis.— Wells & Bryce, 1993:50-51 (photo: species 49)
Tyrodina corticalis.— Coleman, 2001: 135 (photo)

Type locality. South Channel, Port Phillip, Victoria, 14-34 m

Distribution. Q, NSW, V, T, SA, WA: 0-100 m

Family Umbraculidae Dall, 1889

See Valdés (2001b) and Willan & Burn (2003) for commentaries concerning publication date, authorship and type species of both *Umbraculum* and *Tyrodina*.

***Umbraculum* Schumacher, 1817**

Umbraculum Schumacher, 1817: 55, 177

Type species. *Umbrella chinensis* Lamarck, 1801 = *Patella umbraculum* [Lightfoot] 1786

***Umbraculum umbraculum* (Lightfoot, 1786)**

Patella umbraculum [Lightfoot] 1786: 178
Patella sinicum Gmelin, 1791: 3720
Umbraculum sinicum.— Gabriel, 1962: 201
Umbraculum sinicum.— Wells & Bryce, 1993: 51 (photo: species 50), 52
Umbraculum umbraculum.— Coleman, 2001: 135 (photo)

Type locality. Chinese Seas

Distribution. Circum-global tropical and temperate seas, occurring in waters to 100 m deep off south-eastern Tasmania

Order Pleurobranchida

Superfamily Pleurobranchioidea

Now grouped with the Nudibranchia in the clade Nudipleura (Wägale & Willan, 2000), more recently with the doridiniian Anthobranchia in the new subclade Pleuroanthobranchia (Grande et al., 2004) within the Nudipleura.

Family Pleurobranchidae Gray, 1827

***Berthella* Blainville, 1824**

Berthella Blainville, 1824: 262

Type species. *Berthella porosa* Blainville, 1824 = *Bulla plumula* Montagu, 1803

***Berthella medietas* Burn, 1962**

Berthella medietas Burn, 1962b: 142, pl. 1, fig 3; pl. 2, figs 7-8
Berthella medietas.— Burn, 1989: pl. 48.2 (photo)

Type locality. Flinders, Victoria

Distribution. NSW, V, T, SA, WA, NZ: 0-50 m

***Berthella serenitas* Burn, 1962**

Berthella serenitas Burn, 1962b: 143-144, pl. 1, fig 4; pl. 2, figs 5-6

Type locality. Flinders, Victoria

Distribution. V: 0 m

***Berthellina* Gardiner, 1936**

Berthellina Gardiner, 1936: 198

Type species. *Berthellina engeli* Gardiner, 1936

***Berthellina citrina* (Rüppell & Leuckart, 1828)**

- Pleurobranchus citrina* Rüppell & Leuckart, 1828: 20, pl. 1, fig. 1
Pleurobranchus punctatus Quoy & Gaimard, 1832: 299, pl. 22, fig. 14
Berthellina citrina.— Burn, 1989: pl.48.1 (photo)
Berthellina citrina.— Wells & Bryce, 1993: 54, 55 (photo: species 55)
Berthellina citrina.— Edgar, 1997: 276 (photo)
Berthellina citrina.— Coleman, 2001: 136 (photo)

Type locality. Suez, Egypt

Distribution. Circum-global tropical and temperate seas: 0-150 m

Gosliner (2006) states that *Berthellina delicata* (Pease, 1861) “is the common member of the genus found in the western Pacific. There are no external characteristics that distinguish species of *Berthellina*.” Species are separated by differences in “their genitalia”. Identification of south-eastern Australian material will need to be revised.

***Pleurobranchaea* Leue, 1813**

Pleurobranchaea Leue, 1813: 11

Type species. *Pleurobranchaea meckelii* Meckel in Leue, 1813

***Pleurobranchaea maculata* (Quoy & Gaimard, 1832)**

- Pleurobranchidium maculata* Quoy & Gaimard, 1832: 301
Pleurobranchaea maculata.— Burn, 1989: pl.48.3 (photo)
Pleurobranchaea maculata.— Edgar, 1997: 276 (photo)
Pleurobranchaea maculata.— Coleman, 2001: 138 (photo)

Type locality. Southern Australia

Distribution. Q, NSW, V, T, SA, WA, NZ: 0-50 m

***Pleurobranchaea* sp**

Distribution. V: 80-120 m

A small (5 mm long) deep-water species from eastern Bass Strait, in which a shell is retained within a small posterior mantle cavity.

***Pleurobranchus* Cuvier, 1804**

Pleurobranchus Cuvier, 1804: 275

Type species. *Pleurobranchus peronii* Cuvier, 1804

***Pleurobranchus hilli* (Hedley, 1894)**

- Oscanius hilli* Hedley, 1894: 127
Pleurobranchus hilli.— Burn, 1989: pl.47.6 (photo)
Pleurobranchus ovalis.— Burn, 1990: 10 (*non* Pease, 1860)

Type locality. Off Sow & Pigs Reef, Port Jackson, NSW, 17 m

Distribution. NSW, V, T, SA: 0-46 m

The record of *Pleurobranchus ovalis* Pease, 1860 from San Remo, Westernport (Burn, 1990:10) is based upon a 10 mm long juvenile specimen of this large (to 400 mm long) species.

Order Pteropoda

The pteropods of this checklist include the thecosomes as listed by Macpherson & Gabriel (1962), plus an additional species recently collected alive at Wilsons Promontory, and several others the distribution of which (van der Spoel, 1967, 1976) may include eastern Victoria and the Bass Strait area. Gymnosomes have not been reported from Victorian waters. Those listed below have distributions that include eastern and

south-eastern Australia, and may at times extend their range into local waters. Newman (1998) gives an overview of the Australian fauna and its distribution.

Klussmann-Kolb & Dinapoli (2006) review the systematic position of both the Thecosomata and Gymnosomata, showing these to be “sister groups and together closely related to Anaspidea.”

Suborder Thecosomata**Family Limacinidae Gray, 1840*****Limacina* Bosc, 1817**

- Limacina* Bosc, 1817: 42
Spiratella Blainville, 1817: 407

Type species. *Clio helicina* Phipps, 1774

****Limacina bulimoides* (d’Orbigny, 1836)**

Atlanta bulimoides d’Orbigny, 1836: 179

Type locality. Atlantic Ocean

Distribution. Eastern Victoria

****Limacina helicina* (Phipps, 1774)**

Clio helicina Phipps, 1774: 195

Type locality. Northern Atlantic Ocean

Distribution. van der Spoel (1967) maps this species to include the southern half of Tasmania

****Limacina helicoides* Jeffreys, 1877**

Limacina helicoides Jeffreys, 1877: 338

Type locality. North Atlantic Ocean

Distribution. van der Spoel (1967) plots a distributional centre off the central coast of New South Wales

****Limacina inflata* (d’Orbigny, 1836)**

Atlanta inflata d’Orbigny, 1836: 174

Type locality. Mid-Atlantic Ocean

Distribution. Eastern Victoria

****Limacina leseuri* (d’Orbigny, 1836)**

Atlanta leseuri d’Orbigny, 1836

Type locality. Atlantic Ocean

Distribution. Eastern Victoria

****Limacina retroversa* (Fleming, 1823)**

Fusus retroversa Fleming, 1823: 498, pl. 15, fig. 2

Type locality. Atlantic Ocean

Distribution. Eastern Victoria

****Limacina trochiformis* (d’Orbigny, 1836)**

Limacina trochiformis d’Orbigny, 1836: 177

Type locality. Atlantic Ocean

Distribution. Eastern Victoria

Family Cavoliniidae Gray, 1850***Cavolinia* Abildgaard, 1791***Cavolinia* Abildgaard, 1791: 175*Type species.* *Anomia tridentata* Niebuhr, 1775 (ex Forsskålms)****Cavolinia gibbosa* (d'Orbigny, 1836)***Hyalaea gibbosa* d'Orbigny, 1836: 95*Type locality.* Atlantic Ocean*Distribution.* Eastern Victoria***Clio* Linnaeus, 1767***Clio* Linnaeus, 1767: 1094*Type species.* *Clio pyramidata* Linnaeus, 1767****Clio pyramidata* Linnaeus, 1767***Clio pyramidata* Linnaeus, 1767: 1094*Type locality.* Atlantic Ocean*Distribution.* Off Melbourne (Pelseneer, 1888)***Creseis* Rang, 1828***Creseis* Rang, 1828a: 305*Type species.* *Creseis virgula* Rang, 1828****Creseis virgula* Rang, 1828***Creseis virgula* Rang, 1828a: 316*Type locality.* Atlantic Ocean*Distribution.* Eastern Victoria***Cuvierina* Boas, 1886***Cuvierina* Boas, 1886: 131*Type species.* *Cuvierina columella* Rang, 1828***Cuvierina columella* (Rang, 1827)***Cuvierina columella* Rang, 1827: 323*Cuvierina minor* McCoy, 1885: 15*Type locality.* Indian Ocean*Distribution.* Portland, Victoria; world seas

When describing a specimen of basking shark caught off Portland, western Victoria, McCoy (1885) gave the name *Cuvierina minor* to the minute "fusiform, pointed and slightly arched at the posterior end, mouth contracted, oblique" shells filling the intestines with a red pulpy mass.

Diacavolinia* van der Spoel, 1987Diacavolinia* van der Spoel, 1987: 78*Type species.* *Hyalaea longirostris* Blainville, 1821****Diacavolinia longirostris* (Blainville, 1821)***Hyalaea longirostris* Blainville, 1821: 81*Type locality.* Atlantic Ocean*Distribution.* Eastern Victoria***Diacria* Gray, 1847***Diacria* Gray, 1847: 203*Type species.* *Hyalaea trispinosa* Blainville, 1821****Diacria trispinosa* (Blainville, 1821)***Hyalaea trispinosa* Blainville, 1821: 82*Type locality.* West Indian Seas*Distribution.* Eastern Victoria***Styliola* Gray, 1850***Styliola* Gray, 1850a: 16*Type species.* *Cleodora subula* Quoy & Gaimard, 1827****Styliola subula* (Quoy & Gaimard, 1827)***Cleodora subula* Quoy & Gaimard, 1827: 233*Styliola recta* (Lesueur ms.) Blainville, 1827: 655*Type locality.* Teneriffe*Distribution.* Off Melbourne (Pelseneer, 1888)**Family Peraclididae Tesch, 1913*****Peraclis* Forbes, 1844***Peraclis* Forbes, 1844: 186*Type species.* *Atlanta reticulata* d'Orbigny, 1836****Peraclis reticulata* (d'Orbigny, 1836)***Atlanta reticulata* d'Orbigny, 1836: 178*Type locality.* Atlantic Ocean*Distribution.* van der Spoel (1976) shows a strong population centre along the central and northern NSW coast and into the Tasman Sea.****Peraclis valdiviae* (Meisenheimer, 1905)***Procybulia valdiviae* Meisenheimer, 1905: 14*Type locality.* Southern Indian Ocean*Distribution.* van der Spoel (1976) indicates a population centre along the central and northern NSW coast. Two other species with population centres further offshore into the Tasman Sea are: *Perclis apicifulva* Meisenheimer, 1906 and *P. moluccensis* Tesch, 1903.**Family Cymbuliidae Gray, 1840*****Corolla* Dall, 1871***Corolla* Dall, 1871: 137*Type species.* *Corolla spectabilis* Dall, 1871***Corolla ovata* (Quoy & Gaimard, 1832)***Cymbulia ovata* Quoy & Gaimard, 1832: 373*Corolla ovata*.—Newman, 1998: 984, fig.16.64A*Type locality.* Ambon, Indonesia*Distribution.* Gulf of Carpentaria, seas north of New Guinea, V.

A new record of Victorian waters: 1 specimen swimming in water column, Gulch/Landing Station area, south-east corner of Wilsons Promontory, 7 April 2000, leg. Glenys Greenwood (MV F110341).

Suborder Gymnosomata

Family Pneumodermatidae Latreille, 1825

***Pneumodermopsis* Keferstein, 1862**

Pneumodermopsis Keferstein, 1862: 645

Type species. *Pneumodermon ciliatum* Gegenbaur, 1855

****Pneumodermopsis (Crucibranchaea) macrochira* Meisenheimer, 1905**

Pneumodermopsis (Crucibranchaea) macrochira Meisenheimer, 1905: 47

Type locality. Indian Ocean

Distribution. van der Spoel (1976) indicates a strong population centre along the NSW coast.

Family Notobranchaeidae Pelseneer, 1886

***Notobranchaea* Pelseneer, 1886**

Notobranchaea Pelseneer, 1886: 224

Type species. *Notobranchaea macdonaldi* Pelseneer, 1886

****Notobranchaea inopinata* Pelseneer, 1887**

Notobranchaea inopinata Pelseneer, 1887: 40

Type locality. east of Japan

Distribution. van der Spoel (1976) shows a population centre along the central to northern NSW coast.

Family Cliopsidae O.G. Costa, 1873

***Cliopsis* Troschel, 1854**

Cliopsis Troschel, 1854: 222

Type species. *Cliopsis krohni* Troschel, 1854

****Cliopsis krohni* Troschel, 1854**

Cliopsis krohni Troschel, 1854: 222

Type locality. Messina, Italy

Distribution. van der Spoel (1976) indicates a strong population centre along the central to northern NSW coast and into the Tasman Sea.

***Pruvotella* Pruvot-Fol, 1932**

Pruvotella Pruvot-Fol, 1932: 511

Type species. *Pneumodermon pellucidus* Quoy & Gaimard, 1833

****Pruvotella danae* Pruvot-Fol, 1942**

Pruvotella danae Pruvot-Fol, 1942: 17

Type locality. Northern Tasman Sea

Distribution. van der Spoel (1976) shows a population centre off the northern NSW coast.

Family Clionidae Rafinesque, 1815

***Thliptodon* Boas, 1886**

Thliptodon Boas, 1886: 174

Type species. *Thliptodon gegenbauri* Boas, 1886

****Thliptodon antarcticus* Meisenheimer, 1906**

Thliptodon antarcticus Meisenheimer, 1906: 144

Type locality. Antarctica

Distribution. van der Spoel (1976) indicated a population centre along the central to northern NSW coast.

****Thliptodon diaphanus* (Meisenheimer, 1903)**

Pteroceanis diaphanus Meisenheimer, 1903: 93

Type locality. Central Pacific Ocean

Distribution. van der Spoel (1976) indicated a population centre along the central to northern NSW coast.

****Thliptodon gegenbauri* Boas, 1886**

Thliptodon gegenbauri Boas, 1886: 174

Type locality. Messina, Italy

Distribution. van der Spoel (1976) indicated a population centre along the central to northern NSW coast.

Family Hydromylidae Pruvot-Fol, 1942 (1862)

***Hydromyles* Gistel, 1848**

Hydromyles Gistel, 1848: 9

Type species. *Psyche globulosa* Rang, 1825

****Hydromyles globulosa* (Rang, 1825)**

Psyche globulosa Rang, 1825: 284

Type locality. Atlantic Ocean

Distribution. van der Spoel (1976) indicated a strong population centre along the NSW coast and into the Tasman Sea.

Order Nudibranchia

Now grouped with the Pleurobranchoidea in the clade Nudipleura (Wägele & Willan, 2000). More recently, it is the Doridina (= Anthobranchia) only that is grouped with the Pleurobranchoidea in the new subclade Pleuroanthobranchia (Grande et al., 2004) within the Nudipleura.

Suborder Doridina

(= Anthobranchia)

Superfamily Onchidoridoidea

(= Anadoridoidea,

= Phanerobranchia)

Family Goniodorididae H. & A. Adams, 1854

***Ancula* Lovén, 1846**

Ancula Lovén, 1846: 137

Type species. *Polycera cristata* Alder, 1841 = *Tritonia gibbosa* Risso, 1818

***Ancula mapae* (Burn, 1961)**

Drepaniella mapae Burn, 1961b: 102-104, with 1 text fig

Eucrairia mapae.—Burn, 1961d: 51

Ancula mapae.—Burn, 1990: 12

Type locality. Point Danger, Torquay, Victoria

Distribution. NSW, V: 0-10 m

***Goniodoridiella* Pruvot-Fol, 1933**

Goniodoridiella Pruvot-Fol, 1933: 116

Type species. *Goniodoridiella savignyi* Pruvot-Fol, 1933

Goniodoridiella savignyi Pruvot-Fol, 1933

Goniodoridiella savignyi Pruvot-Fol, 1933: 117-118, pl. 2, figs 23-26
Goniodoridiella savignyi.— Rudman, 1998: 992
Goniodoridiella savignyi.— Marshall & Willan, 1999: 215, fig 106

Type locality. Gulf of Suez, Egypt

Distribution. Q, NSW, V: 0-10 m

A small Indo-Pacific tropical and temperate species occasionally encountered in Victorian intertidal waters. Feeds on arborescent bryozoans in eastern Australia (Rudman, 1998).

Goniodoris Forbes & Goodsir, 1839

Goniodoris Forbes & Goodsir, 1839: 647

Type species. *Doris nodosa* Montagu, 1808

Goniodoris meracula Burn, 1958

Goniodoris meracula Burn, 1958: 10, pl. 2, fig 10-11

Type locality. Point Danger, Torquay, Victoria

Distribution. V: 0-10 m

Goniodoris sp

Goniodoris meracula.— Burn, 1973a: 204, fig. 12-13 (photo)
Goniodoris meracula.— Burn, 1989: pl.55.4 (photo)

Distribution. NSW, V, T, SA: 0-124 m

Goniodoris meracula is an uncommonly seen largish species (20 mm long) originally and subsequently found nestled into and eating encrusting ascidians. It has wide thin edges of the foot to line the cavity created. *Goniodoris* sp. is a more common smaller compact species (less than 10 mm long) with thick foot edges. It has been found crawling among algae in rock pools, and in deeper-water dredged samples.

Lophodoris G.O.Sars, 1878

Lophodoris G.O.Sars, 1878: 364

Type species. *Goniodoris danielsseni* Friele & Hansen, 1876

Lophodoris sp

Distribution. V, T: 200 m

Body with wide, highly spiculate inverted umbrella-like mantle, and median crest between rhinophores and gills.

Okenia Menke, 1830

Okenia Menke, 1830: 20

Type species. *Idalia elegans* Leuckart, 1828

Okenia mija Burn 1967

Okenia mija Burn, 1967a: 55, figs 4-5

Type locality. Point Danger, Torquay, Victoria

Distribution. NSW, V, T: 0-15 m

Distinguished by the presence of a single papilla in the mid-line behind the gills. Victorian specimens are small (<5 mm) and rare. Central NSW and south-eastern Tasmanian specimens are much larger (14 & 20 mm, 12 mm maximum length respectively) (Coleman, 2001: 52, Sydney *Okenia*; Rudman, 2004) and maybe represent a different species.

Okenia plana Baba, 1960

Okenia plana Baba, 1960: 80-81, pl. 7, figs. 2a-b
Okenia plana.— Rudman, 2004: figs 20C, 29F

Type locality. Toba, Japan

Distribution. Q, NSW, V, tropical and temperate Indo-Pacific: 0-3 m

Okenia sp 1

Okenia mija.— Rudman, 2004: 29 (*non* Burn, 1967)

Distribution. NSW, V, T: 0-130 m

A small (<4 mm) “rubbery” translucent white species with a slender bundle of long spicules visible within the stiff papillae, and with many long spicules visible within the body wall. Differs from the small species reported from South Australia as *Okenia zoobotryon* (Smallwood, 1910) (Rudman, 2004) by the median notch in the cup-like rhinophoral lamellae.

Okenia sp 2

Distribution. V: 0 m

A small (<4 mm) soft creamy white species with no spicules in the flexible papillae and relatively fewer shorter curved spicules in the body wall. Similar to *Okenia mija* but without the single papilla in the mid-line behind the gills.

Okenia sp 3

Distribution. V: 0-55 m

A stout bodied white species with numerous rather spatulate papillae along the pallial margin.

Okenia sp 4

Distribution. V: 10-74 m

A *Goniodoris*-like species with broad foot, high body and narrow dorsum, with several long slender papillae along each side. Smaller and somewhat similar to the North Atlantic *Okenia aspersa* (Alder & Hancock, 1845, ?= *O. quadricornis* (Montagu, 1815)).

Trapania Pruvot-Fol, 1931

Trapania Pruvot-Fol, 1931: 309

Type species. *Drepania fusca* Lafont, 1874

Rudman (1987b) reviewed the genus *Trapania*.

Trapania aureopunctata Rudman, 1987

Trapania aureopunctata Rudman, 1987b: 203

Type locality. Clovelly, Sydney, NSW, 10-15 m

Distribution. NSW, V: 3-15 m

Trapania benni Rudman, 1987

Trapania benni Rudman, 1987b: 193

Trapania benni.— Coleman, 2001: 53 (photo)

Type locality. Clovelly, Sydney, NSW, 10-15 m

Distribution. NSW, V, SA: 0-23 m

Trapania brunnea Rudman, 1987

Trapania brunnea Rudman, 1987b: 190

Trapania brunnea.— Coleman, 2001: 53 (photo)

Type locality. Clovelly, Sydney, NSW, 10-15 m

Distribution. Q, NSW, V, T, SA, WA, NZ: 0-20 m

Family Onchidorididae Gray, 1827

***Acanthodoris* J.E. Gray in M.E.Gray, 1850**

Acanthodoris J.E. Gray in M.E.Gray, 1850b: 103

Type species. *Doris pilosa* Abildgaard in Müller, 1789

***Acanthodoris metulifera* Bergh, 1905**

Acanthodoris metulifera Bergh, 1905: 98-100, pl. 7, figs. 3-6

Type locality. North-West Coast, Tasmania

Distribution. V, T: 0-43 m

This species may be synonymous with the New Zealand *A. mollicella* Abraham, 1877.

***Acanthodoris nanega* Burn, 1969**

Acanthodoris nanega Burn, 1969: 91-92, figs 41-43

Type locality. Point Lonsdale, Victoria

Distribution. V, SA: 0-3.6 m

Fahey & Valdés (2005) present morphological data for this species in their recent review of *Acanthodoris*.

***Corambe* Bergh, 1869**

Corambe Bergh, 1869: 359 (footnote)

Type species. *Corambe sargassicola* Bergh, 1871

Millen & Martynov, 2005 have advanced convincing evidence for inclusion of *Corambe* and associated genera within the family Onchidorididae.

***Corambe* sp**

Doridella sp.— Rudman, 1998: 990, fig 16.71, pl.35.8

Distribution. NSW, V, T: 0-17 m

A very small (<5 m), cryptic species periodically common upon its food, the encrusting bryozoan *Electra pilosa*.

***Diaphorodoris* Iredale & O'Donoghue, 1923**

Diaphorodoris Iredale & O'Donoghue, 1923: 97, 221

Type species. *Doris luteocincta* M. Sars, 1870

***Diaphorodoris* sp**

Distribution. V, T: 0-3 m

A small creamy–yellow species with red margin to the papillate notum.

***Onchidoris* Blainville, 1816**

Onchidoris Blainville, 1816: 96-97

Type species. *Onchidoris leachii* Blainville, 1816

***Onchidoris maugeansis* (Burn, 1958)**

Lamellidoris maugeansis Burn, 1958: 26, pl. 7, figs. 8, 9, text fig. 4

Onchidoris maugeansis.— Coleman, 2001: 53 (photo)

Type locality. Torquay, Victoria

Distribution. V, T: 0-3 m

Family Polyceridae Alder & Hancock, 1845

***Crimora* Alder & Hancock, 1862**

Crimora Alder & Hancock, 1862: 263

Type species. *Crimora papillata* Alder & Hancock, 1862

***Crimora multidigitalis* (Burn, 1957)**

Euphurus multidigitalis Burn, 1957b: 15-16, pl. 2, figs. 1-6

Type locality. [Point Danger], Torquay, Victoria

Distribution. Q, NSW, V, T: 0-55 m

***Kaloplocamus* Bergh, 1879**

Kaloplocamus Bergh, 1879a: 623, footnote

Type species. *Euplocamus croceus* Philippi, 1836 = *Doris ramosa* Cantraine, 1835

***Kaloplocamus ramosus* (Cantraine, 1835)**

Doris ramosus Cantraine, 1835: 383

Kaloplocamus ramosus.— Burn, 1989: pl.55.1 (photo)

Type locality. Adriatic Sea

Distribution. NSW, V, T: 0-130 m

***Polycera* Cuvier, 1817**

Polycera Cuvier, 1817: 390

Type species. *Doris quadrilineata* Müller, 1776

***Polycera hedgpethi* Er. Marcus, 1964**

Polycera hedgpethi Er. Marcus, 1964: 128-131 with 4 text figs

Polycera hedgpethi.— Coleman, 2001: 47 (photo)

Polycera hedgpethi.— Wells & Bryce, 1993: 76, 77 (photo: species 82)

Type locality. Tomales Bay, California, USA

Distribution. NSW, V, SA, WA, NZ, California, Japan, southern Europe, southern Africa: 0-10 m

A member of the international fouling community, first found in California, now common across southern Australia.

***Polycera janjukia* Burn, 1962**

Polycera janjukia Burn, 1962c: 99-100, figs 3-4

Polycera janjukia.— Coleman, 2001: 47 (photo)

Polycera janjukia.— Miller, 2005: 52, pl. 1, fig. 3

Type locality. [Point Danger], Torquay, Victoria

Distribution. NSW, V: 0-20 m

***Polycera melanosticta* Miller, 1996**

Polycera melanosticta Miller, 1996: 444

Polycera melanostricta.— Coleman, 2001: 47 (photo)

Type locality. Devonport Naval Base, Waitemata Harbour, New Zealand

Distribution. NSW, V, NZ: 0-25 m

***Polycera parvula* (Burn, 1958)**

Palio parvula Burn, 1958: 23-24, pl. 6, figs. 2, 3, text fig. 2

Polycera parvula.— Coleman, 2001: 47 (photo)

Polycera parvula.— Miller, 2005: 52, pl. 1, fig. 4

Type locality. [Point Danger], Torquay, Victoria

Distribution. NSW, V, SA: 0-130 m

***Polycera* sp 1**

Polycera sp.— Coleman, 2001: 47 (photo: 'Gaudy Polycera')

Distribution. V: 0-12 m

Similar to *Polycera parvula*, but with a cirlet of small dark-tipped gills.

***Polycera* sp 2**

Polycera sp.— Coleman, 2001: 47 (photo: 'Remarkable *Polycera*')

Distribution. NSW, V: 0-20 m

Similar to *Polycera parvula*; body smooth or slightly papillose, white overlaid with patchy to dense deep pink pigment, gills small, gill appendages smooth, bulbous, orange tipped.

***Polycera* sp 3**

Distribution. V: 8 m

Similar to *Polycera parvula*; pinkish-orange body papillose all over including orange-tipped scoop-like appendages each side of large gills.

***Polycera* sp 4**

Polycera sp.— Coleman, 2001: 47 (photo: 'Portsea *Polycera*')

Distribution. V, SA: 0-10 m

A dark greyish species with white frosting all over, body pustulose, anterior edge of foot with stout tentaculiform corners.

***Polycera* sp 5**

Polycera sp.— Coleman, 2001: 47 (photo: 'Yellow-Speckled *Polycera*')

Distribution. V: 0-10 m

Semitranslucent body with dense yellow speckling all over, gills large.

***Polycera* sp 6**

Distribution. V: 0 m

Similar to *Polycera parvula*, but with dirty pinkish body covered with large yellow spots, each ringed by a white circle.

***Polycera* sp 7**

Distribution. V, SA: 0 m

Similar to *Polycera parvula*, but with body mottled green and yellow.

***Polycera* sp 8**

Distribution. V, T: 50 m

A deep-water species with a single appendage each side of the gills and another on the outer side of each rhinophore. Possibly wrongly assigned to genus and family.

***Polycera* sp 9**

Distribution. V, T: 35 m

A deep-water species with smooth body.

***Tambja* Burn, 1962**

Tambja Burn, 1962c: 98

Type species. *Nembrotha verconis* Basedow & Hedley, 1905

***Tambja verconis* (Basedow & Hedley, 1905)**

Nembrotha (?) *verconis* Basedow & Hedley, 1905: 158, pl. 2, fig. 1-3

Tambja verconis.— Burn, 1989: pl.54.6 (photo)

Tambja verconis.— Edgar, 1997: 277 (photo)

Tambja verconis.— Coleman, 2001: 46 (photo)

Type locality. Off Newland Head, Backstairs Passage, South Australia, 45 m

Distribution. NSW, V, T, SA, WA, NZ: 0-50 m

***Tambja* sp 1**

Tambja sp.— Coleman, 2001: 46 (photo: 'Southern *Tambja*')

Distribution. NSW, V, T, SA: 0-20 m

Dull green body with brown rhinophores and gills. May involve more than one species.

***Tambja* sp 2**

Distribution. V: 0 m

Bright orange body with purple-tipped rhinophores and gills. Body-coloured, rather than yellow, gills separate this species from the tropical species figured by Coleman (2001: 46) as "Orange *Tambja*" from Lord Howe Island, and by Nakano (2004:110) as *Tambja* sp 2 from Japan. Also similar to the tropical Indo-Pacific and northern Australian (Q, NSW, WA) *Tambja limaciformis* (Eliot, 1908), but lacks the white spotting of that species (Marshall & Willan, 1999: 56, fig. 88).

***Thecacera* Fleming, 1828**

Thecacera Fleming, 1828: 283

Type species. *Doris pennigera* Montagu, 1815

***Thecacera pennigera* (Montagu, 1815)**

Doris pennigera Montagu, 1815: 17, pl. 4, fig. 5

Thecacera pennigera.— Willan & Coleman, 1984: 15, fig. 24

Thecacera pennigera.— Rudman, 1998: 993, fig 16.74B

Thecacera pennigera.— Coleman, 2001: 47 (photo)

Type locality. Devonshire, England

Distribution. NSW, V, SA, NZ: 5-10 m

A wide spread member of the international fouling community, recently (2003) photographed off Rosebud, Port Phillip.

Family Gymnodorididae Odhner, 1941***Gymnodoris* Stimpson, 1855**

Gymnodoris Stimpson, 1855b: 379

Type species. *Gymnodoris maculata* Stimpson, 1855

***Gymnodoris alba* (Bergh, 1877)**

Trevelyan alba Bergh, 1877c: 443-443, pl. 57, figs. 1-12

Gymnodoris alba.— Wells & Bryce, 1993: 88 (photo: species 100)

Gymnodoris alba.— Coleman, 2001: 49 (photo)

Type locality. Burias Island, Philippines

Distribution. Q, NSW, V, T, SA, WA, NT, tropical Indo-Pacific: 0-30 m

***Gymnodoris* (Burn, 1957)**

Nembrotha arnoldi Burn, 1957b: 16, pl. 2, figs. 13, 14

Gymnodoris arnoldi.— Willan & Coleman, 1984: 17, fig. 32

Gymnodoris arnoldi.— Burn, 1989: pl.55.2 (photo)

Gymnodoris arnoldi.— Coleman, 2001: 49 (photo)

Type locality. [Point Danger], Torquay, Victoria

Distribution. V, T, SA: 0-60 m

***Paliolla* Burn, 1958**

Paliolla Burn, 1958: 7

Type species. *Polycera cooki* Angas, 1864

***Paliolla cooki* (Angas, 1864)**

Polycera cooki Angas, 1864: 58-59, pl. 5, fig. 6
Paliolla cooki.— Burn, 1989: pl.55.3 (photo)
Paliolla sp.— Wells & Bryce, 1993: 88, 90 (photo: species 104)
Palliolla cooki.— Coleman, 2001: 51 (photo)

Type locality. Botany Bay, NSW

Distribution. NSW, V, T, SA, WA, NT: 0-130 m

Family Aegiridae Fischer, 1883***Aegires Lovén, 1844***

Aegires Lovén, 1844: 49

Type species. *Polycera punctilucens* d'Orbigny, 1837

***Aegires exeches* Fahey & Gosliner, 2004**

Aegires exeches Fahey & Gosliner, 2004: 656, figs. 48E, 58-61

Type locality. Hekili Point, Maui, Hawaii

Distribution. Q, NSW, V, T, tropical Pacific: 0-100 m

Family Vayssieryidae Thiele, 1931

Bouchet & Rocroi (2005: 118) point out the priority of Okadaiaida Baba, 1930 over Vayssieryidae Thiele, 1931, and their preference for using the former name. Prevailing use of Vayssieryidae by opisthobranch workers indicate reversal of precedence is the better option.

Vayssierya Risbec, 1928

Vayssierya Risbec, 1928: 289
Okadaia Baba, 1930: 47
Pellibranchus Ralph, 1944: 24

Type species. *Vayssierya caledonica* Risbec, 1928

***Vayssierya caledonica* Risbec, 1928**

Vayssierya caledonica Risbec, 1928: 290-292
Vayssierya caldeonica.— Rudman, 1998: 997, fig 16.80
Vayssierya caldeonica.— Marshall & Willan, 1999: 211, fig 91
Vayssierya caldeonica.— Coleman, 2001: 48 (photo)

Type locality. Orphelinat Bay, Noumea, New Caledonia

Distribution. Q, NSW, V: 0-3 m

In Victoria, found in association with its food, the serpulid polychaete worm *Salmacina dysteri*.

Suborder Doridina

(= **Cryptobranchia**,

= **Eudoridoidea**)

The sequence of family and genus taxa in the section follows that presented by Valdés (2002: 629).

Superfamily Phyllidioidea**Family Dendrodorididae O'Donoghue, 1924*****Dendrodoris Ehrenberg, 1831***

Dendrodoris Ehrenberg, 1831: signature g₂

Type species. *Dendrodoris lugubris* Ehrenberg, 1831

***Dendrodoris albopurpura* Burn, 1957**

Dendrodoris albopurpura Burn, 1957b: 13, pl. 3, fig. 3, 12
Dendrodoris albopurpura.— Burn, 1989: pl. 53.5 (photo)

Dendrodoris albopurpura.— Wells & Bryce, 1993: 140 (photo: species 180)

Type locality. Flinders, Victoria

Distribution. V, T, SA, WA: 0-10 m

Dendrodoris albopurpura is a big soft species with large purplish black spots on the median part of the back (Burn, 1989: pl.53.3; Wells & Bryce, 1993: species 180), not to be confused with the equally big but firm tropical Indo-Pacific species *D. elongata* Baba, 1936 and *D. albobrunnea* Allan, 1933, both of which have small brown spots on the back (Willan & Coleman, 1984: 41, figs 121-122).

***Dendrodoris aurea* (Quoy & Gaimard, 1832)**

Doris aurea Quoy & Gaimard, 1832: 265, pl. 19, figs. 4-7

Type locality. Jervis Bay, NSW

Distribution. NSW, V, SA, WA: 0-10 m

A large all orange-pink species.

***Dendrodoris arborescens* (Collingwood, 1881)**

Doridopsis arborescens Collingwood, 1881: 134, pl. 10, fig. 15-17
Dendrodoris fumata auctt. non Rüppell & Leuckart
Dendrodoris nigra auctt. non Stimpson, 1855
Dendrodoris nigra.— Wells & Bryce, 1993: 141 (photo: species 182)
Dendrodoris nigra.— Coleman, 2001: 87 (photo: AMPI 61)

Type locality. Slut Island, Haitan Straits, China

Distribution. All Australian states, wide spread Indo-Pacific: 0-10 m.

As foreshadowed (Goddard, 2005), larval characteristics separate this rather distinctively coloured species from its congeners (Brodie & Calado, 2006). The report (Freame, 1935) of a purplish black sea-slug with red border from Victorian water refers to this species.

***Dendrodoris gunnamatta* Allan, 1932**

Dendrodoris gunnamatta Allan, 1932: 97-98, pl. 5, figs. 4-7
Dendrodoris gunnamatta.— Coleman, 2001: 87 (photo)

Type locality. Gunnamatta Bay, Port Hacking, NSW

Distribution. Q, NSW, V, NZ: 0-5 m

An occasional visitor from warmer northern waters.

***Dendrodoris maugeana* Burn, 1962**

Dendrodoris maugeana Burn, 1962c: 104, fig. 8
Dendrodoris sp.— Coleman, 2001: 87 (photo: 'Brown-speckled Dendrodoris')
Dendrodoris maugeana.— Brodie, 2005: 38, fig. 1A

Type locality. Flinders, Victoria

Distribution. V, T, WA: 0-15 m

***Dendrodoris nigra* (Stimpson, 1855)**

Doris nigra Stimpson, 1855b: 380
Actinodoris australis Angas, 1864: 49, pl. 4, fig. 8
Dendrodoris melaena Allan, 1932: 98, pl. 7, fig. 11
Dendrodoris nigra.— Burn, 1989: pl. 53.4 (photo)

Type locality. Loo Choo & Kikaisima Islands, Japan

Distribution. Q, NSW, V, WA, NT: 0-26 m

Victorian records (Burn, 1990) are of large brownish animals with indistinct pale orange submarginal band around notum.

***Doriopsilla* Bergh, 1880**

Doriopsilla Bergh, 1880b: 316

Type species. *Doriopsilla areolata* Bergh, 1880

***Doriopsilla carneola* (Angas, 1864)**

Doris carneola Angas, 1864: 48-49, pl. 4, fig. 7

Dendrodoris carneola.— Wells & Bryce, 1993: 140 (photo: species 181)

Doriopsilla carneola.— Coleman, 2001: 88 (photo)

Type locality. Port Jackson, NSW

Distribution. Q, NSW, V, T, SA, WA: 0-22 m

***Doriopsilla miniata* (Alder & Hancock, 1864)**

Doridopsis miniata Alder & Hancock, 1864: 130, pl. 31, figs. 18, 19

Doriopsilla miniata.— Coleman, 2001: 88 (photo)

Type locality. Waltair, India

Distribution. Q, NSW, V, WA, Indo-Pacific: 0-20 m

***Doriopsilla peculiaris* (Abraham, 1877)**

Doris peculiaris Abraham, 1877: 258, pl. 30, figs. 15-17

Doriopsilla peculiaris.— Burn, 1989: pl. 54.3-4 (photo)

Doriopsilla peculiaris.— Edgar, 1997: 281 (photo)

Doriopsilla peculiaris.— Coleman, 2001: 88 (photo)

Type locality. South Australia (Port Lincoln)

Distribution. V, T, SA, WA: 0-30 m

***Doriopsilla* sp 1**

Distribution. V: 50-150 m

A highly spiculate species with strongly pustulose notum; possibly a form of *Doriopsilla areolata* Bergh, 1880 from the Atlantic.

***Doriopsilla* sp 2**

Distribution. V, T: 85 m

A narrowly elongate species with smooth highly spiculate notum.

Superfamily Doridoidea**Family Actinocyklidae O'Donoghue, 1929*****Hallaxa* Eliot, 1909**

Hallaxa Eliot, 1909: 80-90

Type species. *Halla decorata* Bergh, 1878

***Hallaxa michaeli* Gosliner & Johnson, 1994**

Hallaxa indecora.— Burn, 1958: 27, non Bergh, 1905

Hallaxa michaeli Gosliner & Johnson, 1994: 182

Hallaxa michaeli.— Coleman, 2001: 63 (photo)

Type locality. Batemans Bay, NSW

Distribution. NSW, V, T: 0-55 m

***Hallaxa* sp 1**

Distribution. V: 0 m

Distinguished from preceding species by flatter body and orange-mottled patterning.

***Hallaxa* sp 2**

Distribution. V: 3-6 m

Very small (<5 mm) flattened white species with a single brown radial line each side on the notum. Notum highly spiculate.

Family Chromodorididae Bergh, 1891

Rudman (1984) reviewed the genera assigned to this family, almost all of which occur in the Australian region.

***Cadlina* Bergh, 1878**

Cadlina Bergh, 1878: XXIX

Type species. *Doris repanda* Alder & Hancock, 1842 = *Doris laevis* Linnaeus, 1767

***Cadlina tasmanica* Rudman, 1990**

Cadlina tasmanica Rudman, 1990: 304

Type locality. Bicheno, Tasmania

Distribution. V, T: 0-6 m

***Cadlina* sp 1**

Distribution. V, T, SA: 0-130 m

Similar to the south Western Australian *Cadlina nigrobranchiata* Rudman, 1985, but distinguished by the tripartite colour pattern of the rhinophores and the absence of a yellow notal margin.

***Cadlina* sp 2**

Distribution. V: 200 m

Notum nodulose, known only from preserved material.

***Ceratosoma* J. E. Gray in M. E. Gray, 1850**

Ceratosoma J. E. Gray in M. E. Gray, 1850: 105

Type species. *Polycera cornigera* A. Adams & Reeve in A. Adams, 1848 = *Doris trilobata* J. E. Gray, 1827

Rudman (1988) reviewed the species of this genus, including anatomical details and coloured illustrations of the two Victorian species.

***Ceratosoma amoenum* (Cheeseman, 1886)**

Chromodoris amoena Cheeseman, 1886: 137

Chromodoris amoena.— Burn, 1989: pl. 48.5 (photo)

Ceratosoma amoena.— Wells & Bryce, 1993: 129 (photo: species 163)

Ceratosoma amoena.— Edgar, 1997: 278 (photo)

Ceratosoma amoenum.— Coleman, 2001: 64-5 (photo)

Type locality. Whangaroa Harbour, New Zealand

Distribution. NSW, V, T, SA, WA: 3-20 m

***Ceratosoma brevicaudatum* Abraham, 1876**

Ceratosoma brevicaudatum Abraham, 1876: 142-143, pl. 8, fig. 6

Ceratosoma adelaidae Basedow & Hedley, 1905: 156, pl. 10, fig. 3-4

Ceratosoma brevicaudatum.— Burn, 1989: pl. 50.5 (photo)

Ceratosoma brevicaudatum.— Wells & Bryce, 1993: 130 (photo: species 164)

Ceratosoma brevicaudatum.— Edgar, 1997: 279 (photo)

Ceratosoma brevicaudatum.— Coleman, 2001: 65(photo)

Type locality. Abrolhos Islands, Western Australia

Distribution. NSW, V, T, SA, WA: 0-120 m

***Chromodoris* Alder & Hancock, 1855**

Chromodoris Alder & Hancock, 1855: XVII

Type species. *Doris magnifica* Quoy & Gaimard, 1832

***Chromodoris alternata* (Burn, 1957)**

Glossodoris alternata Burn, 1957b: 17, pl. 1, figs. 10-11
Chromodoris alternata.— Burn, 1989: pl. 48.4 (photo)
Chromodoris alternata.— Wells & Bryce, 1993: 124 (photo: species 155)
Chromodoris alternata.— Coleman, 2001: 67 (photo)

Type locality. Portarlington, Port Phillip, Victoria

Distribution. V, T, SA, WA: 0-60 m

***Chromodoris ambigua* Rudman, 1987**

Chromodoris ambiguus Rudman, 1987a: 334
Chromodoris sp.— Burn, 1989: pl. 49.6 (photo)
Chromodoris ambigua.— Coleman, 2001: 67 (photo)

Type locality. Griffiths Point, Port Sorrell, Tasmania

Distribution. V, T, SA: 0-20 m

***Chromodoris epicuria* (Basedow & Hedley, 1905)**

Hypselodoris epicuria Basedow & Hedley, 1905: 153-154, pl. 7, figs. 1-3
Glossodoris victoriae Burn, 1957b: 16, pl. 3, fig. 4
Chromodoris epicuria.— Burn, 1989: pl. 49.1-2 (photo)
Chromodoris epicuria.— Wells & Bryce, 1993: 121 (photo: species 149)
Chromodoris epicuria.— Edgar, 1997: 279 (photo)
Chromodoris epicuria.— Coleman, 2001: 70 (photo)

Type locality. Port Willunga, South Australia

Distribution. V, T, SA, WA: 0-15 m

***Chromodoris multimaculosa* Rudman, 1987**

Chromodoris multimaculosa Rudman, 1987a: 331
Type locality. Horseshoe Reef, Devonport, Tasmania
Distribution. V(?), T: 0-15 m

***Chromodoris tasmaniensis* Bergh, 1905**

Chromodoris tasmaniensis Bergh, 1905: 69-70, pl. 5, figs. 12-15
Chromodoris tasmaniensis.— Burn, 1989: pl. 48.6 (photo)
Chromodoris tasmaniensis.— Edgar, 1997: 279 (photo)
Chromodoris tasmaniensis.— Coleman, 2001: 73-74 (photo)

Type locality. North-West Coast, Tasmania

Distribution. NSW, V, T, SA: 0-25 m

This name may represent a species complex.

***Chromodoris tinctoria* (Rüppell & Leuckart, 1828)**

Doris tinctoria Rüppell & Leuckart, 1828: 32, pl. 6, fig. 4
Chromodoris tinctoria.— Burn, 1989: pl. 49.4 (photo)
Chromodoris tinctoria.— Wells & Bryce, 1993: 119 (photo: speices 146)
Chromodoris tinctoria.— Coleman, 2001: 74(photo)

Type locality. Tor, Egypt

Distribution. Q, NSW, V, SA, WA, Indo-Pacific: 0-30 m

***Chromodoris thompsoni* Rudman, 1983**

Chromodoris thompsoni Rudman, 1983: 131
Chromodoris thompsoni.— Burn, 1989: pl. 49.3 (photo)
Chromodoris thompsoni.— Coleman, 2001: 74 (photo)

Type locality. Wattamolla Bay, Sydney, NSW, 24 m

Distribution. NSW, V: 0-10 m

***Chromodoris* sp**

Chromodoris sp.— Coleman, 2001: 75 (photo: 'Haloed Chromodoris')

Distribution. V, SA, WA: 10-15 m

Digidentis Rudman, 1984

Digidentis Rudman, 1984: 226

Type species. *Glossodoris arbuta* Burn, 1961

***Digidentis arbuta* (Burn, 1961)**

Glossodoris arbuta Burn, 1961c: 55-56, pl. 15
Digidentis arbutus.— Burn, 1989: pl. 51.3 (photo)
Digidentis arbuta.— Coleman, 2001: 76 (photo)

Type locality. Point Danger, Torquay, Victoria

Distribution. V, T, WA: 0-85 m

***Digidentis kulonba* (Burn, 1966)**

Hypselodoris kulonba Burn, 1966a: 191
Glossodoris kulonba.— Burn, 1989: pl. 50.2 (photo)

Type locality. Point Lonsdale, Victoria

Distribution. V, T, SA: 0-20 m

An all white species with creamy-yellow notal margins.

***Digidentis perplexa* (Burn, 1957)**

Glossodoris perplexa Burn, 1957b: 17, pl. 3, fig. 1
Digidentis perplexa.— Burn, 1989: pl. 51.4 (photo)
Digidentis perplexa.— Coleman, 2001: 76 (photo)

Type locality. Torquay, Victoria

Distribution. NSW, V, T: 0-30 m

Hypselodoris Stimpson, 1855

Hypselodoris Stimpson, 1855a: 389

Type species. *Goniodoris ? obscura* Stimpson, 1855

***Hypselodoris bennetti* (Angas, 1864)**

Goniodoris bennetti Angas, 1864: 51-52, pl. 4, fig. 10
Hypselodoris bennetti.— Burn, 1989: pl. 50.6 (photo)
Hypselodoris bennetti.— Edgar, 1997: 280 (photo), 281
Hypselodoris bennetti.— Coleman, 2001: 79 (photo)

Type locality. Port Jackson, NSW

Distribution. Q, NSW, V: 0-20 m

Mexichromis Bertsch, 1977

Mexichromis Bertsch, 1977: 113

Type species. *Chromodoris antonii* Bertsch, 1976

***Mexichromis macropus* Rudman, 1983**

Mexichromis macropus Rudman, 1983: 158
Mexichromis macropus.— Burn, 1989: pl. 51.2 (photo)
Mexichromis macropus.— Coleman, 2001: 83 (photo)

Type locality. Western River Cove, Kangaroo Island, South Australia, 10 m

Distribution. Q, NSW, V, SA, WA: 5-20 m

Noumea Risbec, 1928

Noumea Risbec, 1928: 165

Type species. *Noumea romeri* Risbec, 1928

***Noumea aureopunctata* Rudman, 1987**

Noumea aureopunctata Rudman, 1987a: 315

Type locality. West of Don River mouth, Devonport, Tasmania, 7 m

Distribution. V, T: 7-12 m

Reported from Port Phillip, Victoria by Rudman (2006) (24 June 2002, Sea Slug Forum.
http://www.seaslugforum.net/noum_aure.htm)

***Noumea closeorum* Rudman, 1986**

Noumea closei Rudman, 1986: 391

Type locality. Boat Harbour, Tasmania

Distribution. V, T, SA: 0-10 m

Similar to preceding species, but turns brown or black upon preservation. Being dedicated to husband and wife, the patronym is here corrected to *closeorum*.

***Noumea haliclona* (Burn, 1957)**

Glossodoris haliclona Burn, 1957b: 17, pl. 3, fig. 2
Noumea margaretae Burn, 1966a: 195, fig. 7-8
Noumea cameroni Burn, 1966a: 193, fig. 5-6
Noumea haliclona.— Burn, 1989: pl. 50.1 (photo)
Noumea haliclona.— Coleman, 2001: 83 (photo)

Type locality. Portarlington, Port Phillip, Victoria

Distribution. Q, NSW, V, T, SA, WA: 0-20 m

***Noumea sulphurea* Rudman, 1986**

Noumea sulphurea Rudman, 1986: 384
Noumea sulphurea.— Coleman, 2001: 84 (photo)

Type locality. Jibbon Head, Port Hacking, NSW

Distribution. NSW, V, T, SA, WA: 0-12 m

Verconia Pruvot-Fol, 1931

Verconia Pruvot-Fol, 1931: 310

Type species. *Albania? verconis* Basedow & Hedley, 1905

***Verconia verconis* (Basedow & Hedley, 1905)**

Albania? verconis Basedow & Hedley, 1905: 154, pl. 4, figs. 1-4
Verconia verconis.— Burn, 1989: pl. 50.4 (photo)
Verconia verconis.— Wells & Bryce, 1993: 135, 137 (photo: species 177)
Verconia verconis.— Coleman, 2001: 86 (photo)

Type locality: Antechamber Bay, Kangaroo Island, South Australia, 45m

Distribution. NSW, V, T, SA, WA: 0-12 m

Family Dorididae Rafinesque, 1815

***Aldisa* Bergh, 1878**

Aldisa Bergh, Bergh, 1878: 38

Type species. *Doris zetlandica* Alder & Hancock, 1854

***Aldisa* sp**

Distribution. V: 29 m

A greenish species as preserved, with two large yellowish depressions along the mid-line of the notum.

***Aphelodoris* Bergh, 1879**

Aphelodoris Bergh, 1879c: 108

Type species: *Aphelodoris antillensis* Bergh, 1879

***Aphelodoris berghi* Odhner, 1924**

Aphelodoris berghi Odhner, 1924: 53-54
Aphelodoris berghi.— Burn, 1989: pl. 53.1 (photo)
Aphelodoris berghii.— Coleman, 2001: 53 (photo)

Type locality. North-West Coast, Tasmania

Distribution. V, T, SA: 0-30 m

Swims by flexing the body.

***Aphelodoris rossquicki* Burn, 1966**

Aphelodoris rossquicki Burn, 1966c: 339-341, figs 10-11, 31
Aphelodoris cf. varia AMPI 375.— Coleman, 2001: 54 (photo)

Type locality. Ocean Beach, Flinders, Victoria

Distribution. V: 0-50 m

The species figured in Coleman (2001: 54) under this name does not appear to be correctly identified. AMPI 375 in Coleman (2001:54) appears to be closer.

***Aphelodoris varia* (Abraham, 1877)**

Doris variabilis Angas, 1864: 44, pl. 4, fig. 1, non Kelaart, 1858
Doris varia Abraham, 1877: 209
Aphelodoris varia.— Edgar, 1997: 278 (photo)
Aphelodoris varia.— Coleman, 2001: 54 (photo)

Type locality. Port Jackson, NSW

Distribution. NSW, V, T: 0-20 m

***Aphelodoris* sp 1**

Distribution. V: 0-3 m

A large orange species with sketchy patches of brown on the notum, somewhat akin to AMPI 374 from 5 m, Bass Strait (Coleman 2001: 54).

***Aphelodoris* sp 2**

Distribution. V: 43 m

A small (20 mm) creamy-yellow species with a series of maroon spots in the median area of the notum, and traces of brownish concentric lines submarginally. Swims by strongly flexing the body.

***Archidoris* Bergh, 1878**

Archidoris Bergh, 1878: 616

Type species. *Doris tuberculata* Cuvier, 1804, non Müller, 1778 = *Doris pseudoargus* Rapp, 1827

Genus synonymized with *Doris* (Valdés, 2002).

***Archidoris wellingtonensis* (Abraham, 1877)**

Doris wellingtonensis Abraham, 1877: 259, pl. 29, figs. 27-28
Archidoris wellingtonensis.— Willan & Coleman, 1984: 32, 33, fig.90 (photo)
Archidoris wellingtonensis.— Coleman, 2001: 55 (photo)

Type locality. New Zealand

Distribution. V, T, NZ: 0-50 m

***Doris* Linnaeus, 1758**

Doris Linnaeus, 1758: 653

Type species. *Doris verrucosa* Linnaeus, 1758

***Doris cameroni* (Allan, 1947)**

Archidoris cameroni Allan, 1947: 450, pl. 42, figs. 6, 7
Doris cf. cameroni.— Wells & Bryce, 1993: 105 (photo: species 126)
Doris cf. cameroni.— Coleman, 2001: 56 (photo)

Type locality. Angourie Pool, NSW

Distribution. NSW, V, T, SA, WA: 0-60 m

This name may involve more than one species.

***Doris* sp 1**

Doris sp.— Coleman, 2001: 57 (photo: 'Eastern Doris')

Distribution. V: 0 m

Bright yellow in colour with a few white star-like marks on the notum. This is possibly the species figured in Coleman (2001:57) as "Eastern Doris".

***Doris* sp 2**

Distribution. V: 10-12 m

A small (≈10mm) elongate oval species of low profile with gills close to posterior end of notum, colour pale yellow with transverse brown bar at level of rhinophores and narrow brown notal margin, notal papillae polygonal, flattened on top with one to five small dark integumental spots.

***Doriopsis* Pease, 1860**

Doriopsis Pease, 1860a: 32

Type species. *Doriopsis granulosa* Pease, 1860

Genus synonymized with *Doris* (Valdés, 2002).

***Doriopsis flabellifera* (Cheseman, 1881)**

Doris flabellifera Cheseman, 1881: 222-223

Type locality. Auckland Harbour, New Zealand

Distribution. V, NZ: 0-20 m

Figured in Willan & Coleman (1984: 35, fig. 104).

***Neodoris* Baba, 1938**

Neodoris Baba, 1938: 13

Type species. *Neodoris tricolor* Baba, 1938

Genus synonymized with *Doris* (Valdés, 2002).

***Neodoris chrysoderma* (Angas, 1864)**

Doris chrysoderma Angas, 1864: 46, pl. 4, fig. 3
Praeglisca chrysoderma.— Burn, 1957b: 19, pl. 1, fig. 1-5
Neodoris chrysoderma.— Burn, 1989: pl. 51.6 (photo)
Neodoris chrysoderma.— Wells & Bryce, 1993: 102, 104 (photo: species 124)
Neodoris chrysoderma.— Edgar, 1997: 278 (photo)
Neodoris chrysoderma.— Coleman, 2001: 57 (photo)

Distribution. NSW, V, T, SA, WA: 0-50 m

Family Discodorididae Bergh, 1891

***Alloiodoris* Bergh, 1904**

Alloiodoris Bergh, 1904: 41

Type species. *Alloiodoris marmorata* Bergh, 1904

***Alloiodoris marmorata* Bergh, 1904**

Alloiodoris marmorata Bergh, 1904: 42-44, pl. 3, figs. 12-19
Alloiodoris marmorata.— Coleman, 2001: 53 (photo)

Type locality. North-West Coast, Tasmania

Distribution. V, T, SA: 0-33 m

***Discodoris* Bergh, 1877**

Discodoris Bergh, 1877a: 61

Type species. *Discodoris boholiensis* Bergh, 1877

***Discodoris crawfordi* Burn, 1969**

Discodoris crawfordi Burn, 1969: 84-85, figs 19-24
Discodoris cf. crawfordi.— Wells & Bryce, 1993: 99 (photo: species 116)
Discodoris crawfordi.— Coleman, 2001: 55 (photo)

Type locality. South Channel, Port Phillip, Victoria, 18 m

Distribution. V, T, SA, WA: 5-30 m

***Discodoris palma* Allan, 1933**

Discodoris palma Allan, 1933: 448-449, pl. 56, figs. 11, 12
Discodoris palma.— Willan & Coleman, 1984: 36, 37, fig.111 (photo)
Discodoris palma.— Coleman, 2001: 56 (photo)

Type locality. Pussy Cat Bay, Cape Banks, NSW

Distribution. Q, NSW, V: 0-3 m

***Discodoris paroa* Burn, 1969**

Discodoris paroa Burn, 1969: 86-88, figs 30-34
Discodoris paroa.— Coleman, 2001: 56 (photo)

Type locality. Shoreham, Westernport, Victoria

Distribution. V: 0-48 m

***Discodoris turia* Burn, 1969**

Discodoris turia Burn, 1969: 86, figs 25-29

Type locality. Waratah Bay, Victoria

Distribution. V: 5-15 m

***Discodoris* sp 1**

Discodoris sp.— Coleman, 2001: 56 (photo: 'Dappled Discodoris')

Distribution. V, SA: 10 m

A pink species with darker blotches.

***Discodoris* sp 2**

Distribution. V, T: 20 m

A small (20 mm) lighter and darker grey species, broadly oval in shape, the notum with a thick jelly-like skin of mucous. Found embedded in shallow hollows that it eats in a sponge of similar colour.

***Hoplodoris* Bergh, 1880**

Hoplodoris Bergh, 1880a: 51

Type species. *Hoplodoris desmoparypha* Bergh, 1880 = *Doris grandiflora* Pease, 1860

***Hoplodoris nodulosa* (Angas, 1864)**

Doris nodulosa Angas, 1864: 48, pl. 4, fig. 6
Doris pustulata Abraham, 1877
Homoiodoris novaezelandiae Bergh, 1904
Hoplodoris nodulosa.— Burn, 1989: pl. 52.6 (photo)
Hoplodoris nodulosa.— Wells & Bryce, 1993: 102, 104 (photo: species 125)
Hoplodoris nodulosa.— Coleman, 2001: 57 (photo)

Type locality. Coodgee Bay, Sydney, NSW

Distribution. Q, NSW, V, T, SA, WA, NZ: 0-55 m

Genus and species reviewed and redescribed by Fahey & Gosliner (2003).

Jorunna Bergh, 1876

Jorunna Bergh, 1876: 414

Type species. *Doris johnstoni* Alder & Hancock, 1845 = *Doris tomentosa* Cuvier, 1804

Jorunna hartleyi (Burn, 1958)

Rostanga hartleyi Burn, 1958: 28, pl. 7, figs. 12, 13, text fig. 5

Type locality. Breamlea, Victoria

Distribution. V, T: 0-10 m

Jorunna pantherina (Angas, 1864)

Doris pantherina Angas, 1864: 47-48, pl. 4, fig. 5

Jorunna pantherina.— Wells & Bryce, 1993: 108, 110 (photo: species 134)

Jorunna pantherina.— Coleman, 2001: 59 (photo)

Type locality. Coodgee Bay, Sydney, NSW

Distribution. Q, NSW, V, SA, WA, NZ: 0-5 m

Jorunna sp

Halgerda graphica.— Burn, 1957: 12

Distribution. NSW, V, SA, WA: 0-25 m

Mistakenly identified and reported from Victoria as *Halgerda graphica* Basedow and Hedley, 1905 by Burn (1957).

Paradoris Bergh, 1884

Paradoris Bergh, 1884b: 686

Type species. *Paradoris granulata* Bergh, 1884

= *Discodoris indecora* Bergh, 1881

For taxonomic revision, see Dayrat (2006).

Paradoris dubia Bergh, 1904

Discodoris dubia Bergh, 1904: 50-52, pl. 3, figs. 29, 30; pl. 4, figs 1, 2

Discodoris egena Bergh, 1904: 54, pl. 4, fig. 7-14

Alloiodoris marmorata Basedow & Hedley, 1905: 152, pl. 8, fig. 1-2 (non Bergh, 1904)

Alloiodoris nivosus Burn, 1958: 29, pl. 2, fig. 14

Discodoris dubia.— Burn, 1989: pl. 52.2 (photo)

Paradoris leuca Miller, 1995: 904

Discodoris dubia.— Coleman, 2001: 55-6 (photo)

Type locality. North-West Coast, Tasmania

Distribution. NSW, V, T, SA, WA, NZ: 0-50 m

Dayrat (2005) recently added *Paradoris leuca* Miller, 1995 from southern New Zealand to the synonymy of this species. The writer agrees with Valdés (2001a) that the absence in *Paradoris dubia* of accessory glands and copulatory sacs armed with spines associated with the reproductive organs suggests generic assignment elsewhere.

Platydoris Bergh, 1877

Platydoris Bergh, 1877a: 73

Type species. *Doris argo* Linnaeus, 1767

Genus reviewed by Dorgan, Valdés and Gosliner (2002). The following species is listed with little new data.

Platydoris galbana Burn, 1958

Platydoris galbanus Burn, 1958: 13-14, pl. 1, fig 6-7

Platydoris galbana.— Coleman, 2001: 61 (photo)

Type locality. Sutherlands Bay, Phillip Island, Victoria

Distribution. NSW, V: 0-30 m

Listed as a protected species under the Flora and Fauna Guarantee Act, Victoria.

Rostanga Bergh, 1879

Rostanga Bergh, 1879b: 353

Type species. *Doris concinna* Alder & Hancock, 1848 = *Doris rubra* Risso, 1818

Rudman & Avern (1989) reviewed the genus *Rostanga*, providing colour illustrations of each of the species reported from Victoria and information on their egg ribbons.

Rostanga australis Rudman & Avern, 1989

Rostanga australis Rudman & Avern, 1989: 312

Rostanga australis.— Coleman, 2001: 61 (photo)

Type locality. Portsea Pier, Port Phillip Bay, Victoria

Distribution. V, T, SA, WA: 0-20 m

Rostanga bassia Rudman & Avern, 1989

Rostanga bassia Rudman & Avern, 1989: 310

Type locality. West Head, Flinders, Victoria

Distribution. V: 0 m

Rostanga bifurcata Rudman & Avern, 1989

Rostanga bifurcata Rudman & Avern, 1989: 293

Rostanga bifurcata.— Wells & Bryce, 1993: 108, 109 (photo: species 132)

Type locality. Inscription Point, Kurnell, Sydney, NSW, 12 m

Distribution. Q, NSW, V, WA, tropical Indo-West Pacific: 5-19 m.

A new record for Victorian waters: 1 specimen and egg ribbon, 19 m on sponge, Kessops Cove, east coast of Wilsons Promontory, 2 January 1998, leg. Glenys Greenwood. (MV F110092).

Rostanga calumus Rudman & Avern, 1989

Rostanga calumus Rudman & Avern, 1989: 300

Rostanga arbutus auctt non Angas, 1864

Rostanga calumus.— Wells & Bryce, 1993: 108, 109 (photo: species 133)

Rostanga calumus.— Coleman, 2001: 62 (photo)

Distribution. NSW, V, T, SA, WA: 0-30 m

Type locality. Woolgoolga, NSW

Sclerodoris Eliot, 1904

Sclerodoris Eliot, 1904: 355, 360-361

Type species. *Doris osseosa* Kelaart, 1859

Iredale & McMichael (1962:93) designated *Doris osseosa* Kelaart, 1859 as type species of *Sclerodoris*, and were followed in this by Thompson & Brown (1974). Rudman (1978:84) however selected *Sclerodoris tuberculata* Eliot, 1904 as type

species, and Valdés & Gosliner (2001:166) and Valdés (2001a:296) subsequently re-designated the same species as type. As justification for their action, Valdés & Gosliner (2001) expressed doubt as to the identification of the material identified by Eliot (1904) as *Doris osseosa*.

***Sclerodoris tarka* Burn, 1969**

Sclerodoris tarka Burn, 1969: 88-90, figs 35-40

Type locality. Point Lonsdale, Victoria

Distribution. V, T: 0-43 m

The species figured in Burn (1989: pl 52, fig.4) under this name is *Sclerodoris* sp. below.

***Sclerodoris trenberthi* (Burn, 1962)**

Asteronotus (Tumbia) trenberthi Burn, 1962a: 161-163, pl. 1, figs. 3-5, text figs. 13, 14

Sclerodoris trenberthi.— Burn, 1989: pl. 52.5 (photo)

Type locality. Fiddlers Bay, Spencer Gulf, South Australia

Distribution. V, SA: 0-10 m

***Sclerodoris* sp**

Sclerodoris tarka.— Burn, 1989: pl. 52.4 (photo)

Distribution. V, T, SA: 0-12 m

A brownish-orange species to 40 mm long, notum crowded with small conical papillae interspersed with scattered larger papillae, always with a prebranchial brown patch. Figured in Burn (1989: pl.52.4), where it is misidentified as *Sclerodoris tarka*.

***Thordisa* Bergh, 1877**

Thordisa Bergh, 1877b: 540

Type species. *Thordisa maculigera* Bergh, 1877

***Thordisa sanguinea* Baba, 1955**

Thordisa sanguinea Baba, 1955: 47, pl. 10, figs. 25, 26, text fig. 25

Thordisa sanguinea.— Coleman, 2001: 62 (photo)

Type locality. Sagami Bay, Japan, 10 m

Distribution. Q, NSW, V, Japan: 0-15 m

***Thordisa verrucosa* (Crosse in Angas, 1864)**

Goniodoris verrucosa Crosse in Angas, 1864: 56-57, pl. 5, fig. 4

Thordisa sabulosa Burn, 1957: 20, pl. 1, fig. 6-9

Thordisa verrucosa.— Coleman, 2001: 63 (photo)

Type locality. Shark Island, Port Jackson, NSW

Distribution. Q, NSW, V: 0-65 m

***Trippa* Bergh, 1877**

Trippa Bergh, 1877a: 63

Type species. *Trippa ornata* Bergh, 1877

Valdés & Gosliner (2001:111) synonymized *Trippa* with *Atagema* J. E. Gray, 1850. Their concept of the type species of *Atagema*, *Doris carinata* Quoy & Gaimard, 1832 was based upon examination of the 9 mm long, poorly preserved syntype and portion of its radula, and the brief description and figure of living material identified by Willan and Coleman (1984: species 100) as *Atagema carinata*. Bergh's (1904: 39-41, pl. 3, fig. 8-11) description and figures refer to the same large (to 70 mm) species that Willan and Coleman (1984) had before them, but this species is not necessarily Quoy & Gaimard's (1832) small

species. Bergh's *Atagema carinata* has a very long, winding, slender prostate and very short muscular vas deferens (Burn, pers. obs.).

***Trippa albata* Burn, 1962**

Trippa albata Burn, 1962c: 101, fig 5

Type locality. Sutherlands Bay, Phillip Island, Victoria

Distribution. V, WA: 0-74 m

Suborder Dendronotina

Family Tritoniidae Lamarck, 1809

***Marianina* Pruvot-Fol, 1930**

Marianina Pruvot-Fol, 1930: 229

Type species. *Mariana rosea* Pruvot-Fol, 1930

Marianina is retained in Tritoniidae following Willan (1988), rather than maintaining a separate family for this monotypic genus.

***Marianina rosea* (Pruvot-Fol, 1930)**

Mariana rosea Pruvot-Fol, 1930: 229

Marianina rosea.— Willan & Coleman, 1984: 50, 51, fig. 165 (photo)

Marianina rosea.— Burn, 1989: pl. 43.5 (photo)

Marianina rosea.— Marshall & Willan, 1999: 247, fig. 233 (photo)

Marianina rosea.— Coleman, 2001: 95 (photo)

Type locality. Kuto, New Caledonia

Distribution. Q, NSW, V, SA, WA, tropical and temperate Indo-Pacific: 0-25 m

***Marioniopsis* Odhner, 1934**

Marioniopsis Odhner, 1934: 286

Type species. *Tritonia cyanobranchiata* Rüppell & Leuckart, 1828

Smith & Gosliner (2005) suggest uniting *Marioniopsis* with the earlier *Marionina* Vayssière, 1877.

***Marioniopsis platyctenea* Willan, 1988**

Marioniopsis platyctenea Willan, 1988: 49

Marioniopsis platyctenea.— Wells & Bryce, 1993: 168, 170 (photo: species 219)

Type locality. Julian Rocks, Cape Byron, NSW, 11 m

Distribution. NSW, V, T, WA: 3-20 m

***Paratritonia* Baba, 1949**

Paratritonia Baba, 1949: 84-85, 166

Type species. *Paratritonia lutea* Baba, 1949

***Paratritonia lutea* Baba, 1949**

Paratritonia lutea Baba, 1949: 85-86, 166, pl. 34, figs. 123, text figs. 104-106

Paratritonia cf. lutea 'Sea Fan Tritonia'.— Coleman, 2001: 96 (photo)

Type locality. Sagami Bay, Japan, 60-100 m

Distribution. V, SA, Japan: 3-10 m

Probably needs to be re-identified.

***Tritonia* Cuvier, 1797**

Tritonia Cuvier, 1797: 387

Type species. *Tritonia hombergii* Cuvier, 1803

***Tritonia* sp 1**

Distribution. V: 0 m

Small (<10 mm), pale bluish pink, long velar processes.

***Tritonia* sp 2**

Distribution. V: 0-5 m

Small (<10 mm), fawn to pale brown, velum with projecting corners only, branching of lateral processes ill-formed.

***Tritonia* sp 3**

Tritonia sp.— Coleman, 2001: 97 (photo: 'Patchwork Tritonia')

Tritonia sp.— Coleman, 2001: 97 (photo: 'Carijoa Tritonia')

Distribution. Q, NSW, V, T, SA, WA: 0-30 m

To 30 mm length, orange body with white velar and lateral processes

***Tritonia* sp 4**

Tritonia sp.— Coleman, 2001: 97 (photo: 'Latticed Tritonia')

Distribution. V: 3-10 m

To 25 mm length, white body with paler whitish patches, velar and lateral processes white.

***Tritonia* sp 5**

Distribution. V: 5 m

To 20 mm length, fawn body with brown reticulum on notum, velar and lateral processes brownish.

***Tritonia* sp 6**

Distribution. V: 50 m

To 15 mm length, notum papillose in known preserved material. A deeper water species possibly to be identified with a tropical or subtropical species.

***Tritonia* sp 7**

Distribution. V: 600 m

To 30 mm length, in life body blotched red and fawn.

***Tritonia* sp 8**

Tritonia nilsodhneri.— Coleman, 2001: 97 (photo)

Distribution. NSW, V: 8-20 m

To 25 mm length, body colour orange, notum whitish (NSW) or with pale whitish tracery (V). Figured in Coleman (2001: 97), where it is misidentified as the western European *Tritonia nilsodhneri* Ev. Marcus, 1983.

Family Dendronotidae Allman, 1845***Dendronotus* Alder & Hancock, 1845**

Dendronotus Alder & Hancock, 1845: 47, fam. 3, pl. 3

Type species. *Doris arborescens* Müller, 1776,? = *Amphitrite frondosa* Ascanius, 1774

***Dendronotus* sp**

Distribution. V, T: 10-80 m

Small (<10 mm) pinkish-red species with densely ramose lateral appendages. Known only from subtidal to deep water.

Family Dotidae Gray, 1845***Doto* Oken, 1815**

Doto Oken, 1815: 278

Type species. *Doris coronata* Gmelin, 1791

***Doto ostenta* Burn, 1958**

Doto ostentus Burn, 1958: 16, pl. 1, fig 5, text fig 9

Doto ostenta.— Burn, 1989: pl. 56.5(photo)

Doto ostenta.— Beesely et al., 1998: pl. 36, figs 6-7(photo)

Doto ostenta.— Coleman, 2001: 99(photo)

Type locality. Torquay, Victoria

Distribution. NSW, V, T, SA: 0-85 m

A rare wholly black form may represent another species. This is the only *Doto* species in Victorian waters with a black spot in the tip of the secondary papillae on the cerata.

***Doto pita* Er.Marcus, 1955**

Doto pita Er. Marcus, 1955: 169-170, pl. 24, figs. 161-167

Doto pita.— Coleman, 2001: 99 (photo)

Type locality. São Sebastião, Brazil

Distribution. V, NZ, Japan: 0-95 m

Originally described from Brazil, *Doto pita* has since been reported from many of the world's seas. Cerata irregularly shaped, rather angular.

***Doto* sp 1**

Distribution. V: 0 m

Cerata strongly curved, secondary papillae on outer side only. Similar to the North Atlantic *Doto doerga* Ev. & Er. Marcus, 1963

***Doto* sp 2**

Doto sp.— Coleman, 2001: 99 (photo: 'Red Backed Doto')

Distribution. NSW, V, T: 5-33 m

White body with red stripe along middle of notum branching to each rhinophore, secondary papillae on cerata brownish.

***Doto* sp 3**

Doto sp.— Coleman, 2001: 99 (photo: 'Coleman's Doto')

Distribution. V, T: 10-40 m

White body with pink spots on sides and notum, cerata long, fawn.

***Doto* sp 4**

Distribution. V: 30 m

Cerata orange with black speckling.

Family Scyllaeidae Alder & Hancock, 1855

***Crosslandia* Eliot, 1902**

Crosslandia Eliot, 1902: 64

Type species. *Crosslandia fusca* Eliot, 1902 = *Crosslandia viridis* Eliot, 1902

***Crosslandia viridis* Eliot, 1902**

Crosslandia viridis Eliot, 1902: 64-68, pl. 5, figs 1-8, text figs. 2-4
Crosslandia viridis.— Burn, 1989: pl. 56.3 (photo)
Crosslandia viridis.— Wells & Bryce, 1993: 174 (photo: species 224)
Crosslandia viridis.— Coleman, 2001: 100 (photo)

Type locality. Zanzibar, Africa

Distribution. Q, V, WA, tropical Indo-Pacific: 0-10 m

***Scyllaea* Linnaeus, 1758**

Scyllaea Linnaeus, 1758: 656

Type species. *Scyllaea pelagica* Linnaeus, 1758

***Scyllaea pelagica* Linnaeus, 1758**

Scyllaea pelagica Linnaeus, 1758: 656
Scyllaea pelagica.— Burn, 1989: pl. 56.2 (photo)
Scyllaea pelagica.— Wells & Bryce, 1993: 173 (photo: species 223)
Scyllaea pelagica.— Coleman, 2001: 100 (photo)

Type locality. “in Pelagi Fuco Natante”

Distribution. All tropical and temperate seas: 5-10 m

First reported from Victoria by Hedley (1895).

Family Tethydidae Rafinesque, 1815

***Melibe* Rang, 1829**

Melibe Rang, 1829: 129

Type species. *Melibe rosea* Rang, 1829

***Melibe australis* (Angas, 1864)**

Melibaea australis Angas, 1864: 62-62, pl. 6, fig. 2
Melibe australis.— Burn, 1989: pl. 56.4 (photo)
Melibe australis.— Wells & Bryce, 1993: 174, 175 (photo: species 225)
Melibe australis.— Coleman, 2001: 100 (photo)

Type locality. Watsons Bay, Port Jackson, NSW

Distribution. NSW, V, T, SA, WA: 0-10 m

***Melibe maugeana* Burn, 1960**

Melibe pellucida Burn, 1957b: 24, pl. 3, fig. 5-7, non Bergh, 1904
Melibe maugeana Burn, 1960c: 70

Type locality. Torquay, Victoria

Distribution. V: 0-10 m

Cerata long, round in section, single or double pointed at apex, rhinophore sheath with wing-like keel behind.

***Melibe* sp 1**

Distribution. V: 0 m

Cerata short, swollen, club-shaped, with finely-pointed short papillae all over.

***Melibe* sp 2**

Distribution. V: 0 m

Like *Melibe maugeana*, but with long slender papillae projecting from cerata.

***Melibe* sp 3**

Distribution. V: 0-10 m

Body greenish orange, angular; cerata long, apically wedge-shaped.

Suborder Arminina

Family Arminidae Iredale & O'Donoghue, 1923

***Armina* Rafinesque, 1814**

Armina Rafinesque, 1814b: 30

Type species. *Armina tigrina* Rafinesque, 1814

***Armina* sp 1**

Armina sp.— Edgar, 1997: 283 (photo)
Dermatobranchus sp.— Coleman, 2001: 103 (photo: ‘Coastal Dermatobranchus’)

Distribution. NSW, V, SA, WA: 2-55 m

A periodically common black and white species.

***Armina* sp 2**

Distribution. V: 55 m

Greenish animal with minute dark spots, known from dredged specimens. Probably to be identified with a warmer species occurring in Queensland or New South Wales.

***Dermatobranchus* van Hasselt, 1824**

Dermatobranchus van Hasselt, 1824: 37

Type species. *Dermatobranchus striatus* van Hasselt, 1824

***Dermatobranchus pulcherrimus* Miller & Willan, 1986**

Dermatobranchus pulcherrimus Miller & Willan, 1986: 384-5

Type locality. Colville Channel, Outer Hauraki Gulf, New Zealand, 55 m

Distribution. V, T, NZ: 0-30 m

Animal pinkish-red with slender longitudinal white ridges on notum.

***Heterodoris* Verrill & Emerton, 1882**

Heterodoris Verrill & Emerton, 1882: 548-549

Type species. *Heterodoris robusta* Verrill & Emerton, 1882

***Heterodoris* sp**

Distribution. NSW, V: 600 m

A large (>50 mm) species, pale flesh colour in life.

Family Zephyrinidae Iredale & O'Donoghue, 1923

***Caldukia* Burn & Miller, 1969**

Caldukia Burn & Miller, 1969: 23-24

Type species. *Proctonotus* ? *affinis* Burn, 1958

***Caldukia affinis* (Burn, 1958)**

Proctonotus ? *affinis* Burn, 1958: 32-33, pl. 7, figs. 15, text fig. 8
Caldukia affinis.— Burn, 1989: pl. 56.1 (photo)
Caldukia affinis.— Coleman, 2001: 104 (photo)

Type locality. Torquay, Victoria

Distribution. NSW, V, T, SA: 0-55 m

Janolus Bergh, 1884

Janolus Bergh, 1884a: 18

Type species. *Janolus australis* Bergh, 1884

Janolus hyalinus (Alder & Hancock, 1854)

Antiopa hyalina Alder & Hancock, 1854: 105

Type locality. Hilbro Island, Dee River, England

Distribution. V, NZ: 0-18 m

This species is considered an introduction to south-eastern Australia and New Zealand (Miller & Willan, 1986) from European waters. Reassessment and revision of the species is desirable.

Janolus sp 1

Janolus sp.— Coleman, 2001: 104 (photo: 'Ringed Janolus')

Distribution. NSW, V, T, SA, WA: 0-20 m

An orange species with numerous crowded smooth cerata. Victorian and Tasmanian specimens attain 15-20 mm in length, elsewhere it can exceed 40 mm. May involve more than one species.

Janolus sp 2

Antiopella sp.— Coleman, 2001: 103 (photo: 'Lined Antiopella')

Distribution. V: 0-10 m

A pale species with brown patches on the body and silvery-white lines along the cerata.

Janolus sp 3

Distribution. V: 0-3 m

Differs from preceding species by irregular bands of golden flecks around the cerata.

Family Madrellidae Preston, 1911

Madrella Alder & Hancock, 1864

Madrella Alder & Hancock, 1864: 141-142

Type species. *Madrella ferruginosa* Alder & Hancock, 1864

Madrella sanguinea (Angas, 1864)

Janus sanguineus Angas, 1864: 63-64, pl. 6, fig. 5

Madrella sanguinea.— Burn, 1989: pl. 55.6 (photo)

Madrella sanguinea.— Coleman, 2001: 103 (photo)

Type locality. Watsons Bay, Port Jackson, NSW

Distribution. Q, NSW, V, T, SA, WA: 0-25 m

Lives on, eats, and is same colour as the encrusting bryozoan *Mucropetraliella ellerii*.

Suborder Aeolidina

Family Flabellinidae Bergh, 1889

Flabellina Griffith & Pidgeon, 1833

Flabellina Griffith & Pidgeon, 1833: 40

Type species. *Doris affinis* Gmelin, 1791

Voigt, 1834 is usually cited as author and date for *Flabellina*. An earlier introduction, from 16 December 1833, is available in Part 38 of the English edition by Griffith & Pidgeon of Cuvier's *Le Règne Animal*.

Opinion is still divided over the question of union or separation of *Flabellina* and *Coryphella* J.E. Gray, 1850.

Flabellina poenicia (Burn, 1957)

Hervia poenicia Burn, 1957b: 25, pl. 2, figs. 7-10

Coryphellina poenicia aurantia Burn, 1962: 109

Flabellina poenicia.— Coleman, 2001: 106 (photo)

Distribution. V, T, SA: 0-55 m

Type locality. Breamlea, Victoria

The name of this species was originally mistakenly spelled *peonicia*, and overlooked at proof stage prior to publication. It has been spelled *poenicia* ever since, the Latin name meaning purple-red, in allusion to the colours of the living type material. Article 33.3.1 (ICZN, 1999) permits retention of "incorrect subsequent spelling in prevailing usage".

Flabellina rubrolineata (O'Donoghue, 1929)

Coryphellina rubrolineata O'Donoghue, 1929: 798-802, text fig. 219

Flabellina rubrolineata.— Wells & Bryce, 1993: 151 (photo: species 196)

Flabellina rubrolineata.— Edgar, 1997: 282, 283 (photo)

Flabellina rubrolineata.— Coleman, 2001: 106 (photo)

Type locality. Suez, Egypt

Distribution. Q, NSW, V, WA, NT, Indo-Pacific: 0-30 m

Flabellina sp 1

Flabellina sp.— Coleman, 2001: 106 (photo: 'Orange-tipped Flabellina')

Distribution. V, T: 0-50 m

This species has a purple body with darker extremities, wrinkled rhinophores, and tentaculiform foot corners.

Flabellina sp 2

Distribution. V: 0-10 m

Very similar to the preceding species, differing in the broadly rounded and expanded anterior foot.

Flabellina sp 3

Distribution. V: 10-30 m

Separated from *Flabellina poenicia* and *F. rubrolineata* by the presence of long slender tentaculiform foot-corners and mulberry-like clavus of the rhinophores.

Tularia Burn, 1966

Tularia Burn, 1966e: 26

Type species. *Cuthona bractea* Burn, 1962

Tularia bractea (Burn, 1962)

Cuthona bractea Burn, 1962c: 110-111, text figs. 11-12

Tularia bractea.— Coleman, 2001: 109 (photo)

Type locality. Torquay, Victoria

Distribution. V, T, SA, NZ: 0-15 m

Family Eubranchidae Odhner, 1934***Eubranchus* Forbes, 1838***Eubranchus* Forbes, 1838: 5*Type species. Eubranchus tricolor* Forbes, 1838***Eubranchus rubeolus* Burn, 1964***Eubranchus rubeolus* Burn, 1964b: 13-14, figs 6-10*Type locality.* Point Lonsdale, Victoria*Distribution.* V: 0 m

Specimens recorded from New Zealand under this name (Miller, 1971) need to be re-identified.

Eubranchus* sp 1Distribution.* V: 0-10 m

A brownish species (to 15 mm length) with three circlets of bluntly pointed secondary papillae on the cerata, and the anus opening behind two rows of cerata on the right side.

Eubranchus* sp 2Distribution.* V: 0-10 m

Very similar to preceding species in shape and colour, distinguished by smaller size (7 mm) and anal opening behind three rows of cerata.

Eubranchus* sp 3Distribution.* V: 0 m

A small white species with a red patch on the head and black spots on the cerata.

Eubranchus* sp 4Distribution.* V: 0 m

A small reddish-brown species with very long tentaculiform foot corners.

Eubranchus* sp 5Distribution.* V: 0 mA small species with inflated smooth cerata, much resembling the European *Eubranchus pallidus* (Alder & Hancock, 1842) in shape and colour pattern.***Eubranchus* sp 6***Distribution.* V: 8-10 m

Body white, to 12 mm long, striped dorsally and laterally deep red, cerata spirally twisted, grey with six slightly knobby yellowish rings.

Eubranchus* sp 7Distribution.* V: 10-12 m

A small (<5 mm) species with white body and slightly knobby inflated cerata, with orange digestive gland.

Eubranchus* sp 8Distribution.* V: 6-10 m

A small (<5 mm) species with dense orange speckling all over body, indigo blue tips to oral tentacles and rhinophores, and black digestive gland in elongate, slightly knobby cerata.

Family Aeolidiidae Gray, 1827***Aeolidiella* Bergh, 1867***Aeolidiella* Bergh, 1867: 99*Type species. Eolis alderi* Cocks, 1852 = *Eolida soemmerringii* auctt (ICZN suppressed)***Aeolidiella drusilla* Bergh, 1900***Aeolidiella drusilla* Bergh, 1900: 233-235, pl.20, figs 41-46
Aeolidiella drusilla.— Coleman, 2001: 115 (photo)*Type locality.* French Pass, Cook Strait, New Zealand*Distribution.* V, T, NZ: 0-58 m***Anteaeolidiella* Miller, 2001***Anteaeolidiella* Miller, 2001: 634*Type species. Aeolidiella indica* Bergh, 1888***Anteaeolidiella foulisi* (Angas, 1864)***Aeolis foulisi* (Angas, 1864): 64-65, pl.6, fig 3
Aeolidiella indica Bergh, 1888: 755, pl. 78, fig. 1-2
Aeolidiella takanosimensis Baba, 1930: 122, pl. 4, fig. 5 a-c
Anteaeolidiella indica.— Coleman, 2001: 115 (photo)*Type locality.* Port Jackson, NSW*Distribution.* Q, NSW, V, wide-spread Indo-Pacific: 0-15 mLike *Aeolidiella takanosimensis* Baba, 1930 from Japan, specimens from Victorian waters are opaque white with a series of reddish orange diamond-shaped patches, each with a white centre, along the mid-line of the body. NSW specimens have a pale to bright orange body with or without patches along the mid-line, the latter agreeing closely with the description and figure of *Aeolis foulisi* Angas, 1864. *Anteaeolidiella foulisi* is an earlier name for the wide-spread species currently known as *A. indica* (Bergh, 1888). It has been suggested that the even earlier *Eolis cacaotica* Stimpson, 1855, also from Port Jackson (= Sydney Harbour), may be identical with *A. foulisi* (Burn, 1965).***Baeolidia* Bergh, 1888***Baeolidia* Bergh, 1888: 777*Type species. Baeolidia moebii* Bergh, 1888***Baeolidia australis* (Rudman, 1982)***Spurilla australis* Rudman, 1982: 164
Spurilla australis.— Burn, 1989: pl. 57.5 (photo).— Wells & Bryce, 1993: 154, 155 (photo: species 201)
Baeolidia australis.— Coleman, 2001: 115-6 (photo)*Type locality.* Pilot Beach, near Laurieton, NSW*Distribution.* Q, NSW, V, SA, WA, NZ: 0-12 m***Burnaia* Miller, 2001***Burnaia* Miller, 2001: 659*Type species. Aeolidia helicochorda* Miller, 1988***Burnaia helicochorda* (Miller, 1988)***Aeolidia helicochorda* Miller, 1988: 391*Type locality.* Goat Island Bay, Leigh, New Zealand*Distribution.* NSW, V, T, SA, NZ: 0-10 m***Cerberilla* Bergh, 1873***Cerberilla* Bergh, 1873: 160*Type species. Cerberilla longicirrho* Bergh, 1873

***Cerberilla incola* Burn, 1974**

Cerberilla incola Burn, 1974a: 54-55, figs 11-14

Type locality. Corio Bay, Port Phillip, Victoria

Distribution. NSW, V, T: 0-50 m

***Cerberilla* sp 1**

Distribution. V, SA: 5-30 m

Larger than the preceding species, with two stripes of black along the dorsal surface of each cerata.

***Cerberilla* sp 2**

Distribution. V, T: 20 m

Head with long black oral tentacles, cerata all yellow.

***Spurilla* Bergh, 1864**

Spurilla Bergh, 1864: 205

Type species. *Eolis neopolitana* Delle Chiaje, 1841

***Spurilla macleayi* (Angas, 1864)**

Aeolis macleayi Angas, 1864: 65-66, pl. 6, fig. 4

Spurilla macleayi.— Burn, 1989: pl. 57.6 (photo)

Spurilla macleayi.— Coleman, 2001: 116 (photo)

Type locality. Port Jackson, NSW

Distribution. NSW, V, T, SA: 0-30 m

It is very probable that *Aeolidiella faustina* Bergh, 1900 from New Zealand (Miller, 2001) and Tasmania (Bergh, 1904) is synonymous with *Spurilla macleayi*.

Family Glaucidae Gray, 1827

This family is restricted to the truly pelagic species, rather than embracing the whole of the facelinid aeolids (Willan, 1987; Valdés & Campillo, 2004).

***Glaucus* Forster, 1777**

Glaucus Forster, 1777: 49

Type species. *Glaucus atlanticus* Forster, 1777

The genus *Glaucilla* Bergh, 1860 has now been reduced to synonymy with *Glaucus* (Valdés & Campillo, 2004).

***Glaucus atlanticus* Forster, 1777**

Glaucus atlanticus Forster, 1777: 49

Glaucus atlanticus.— Coleman, 2001: 115 (photo)

Type locality. South Atlantic Ocean

Distribution. Eastern V, world-wide seas: 0 m

Open ocean pelagic species, washed ashore after storms.

***Glaucus marginatus* (Bergh, 1860)**

Glaucilla marginatus Bergh, 1860: 325, pl. 8, fig. 9

Glaucilla marginata.— Coleman, 2001: 115 (photo)

Type locality. not recorded [Hawaiian Islands]

Distribution. Eastern V, Pacific: 0 m

Open ocean pelagic species, washed ashore after storms. Far less common than preceding species.

Family Facelinidae Bergh, 1889***Austraeolis* Burn, 1962**

Austraeolis Burn, 1962c: 120

Type species. *Flabellina ornata* Angas, 1864

***Austraeolis ornata* (Angas, 1864)**

Flabellina ornata Angas, 1864: 67-68, pl.6, fig. 7

Rizzolia australis Bergh, 1884: 27, pl. 9, fig. 1-5

Austraeolis westralis Burn, 1966e: 31, fig. 11-14

Austraeolis ornata.— Wells & Bryce, 1993: 157, 159 (photo: species 206).— Coleman, 2001: 110 (photo)

Type locality. Port Jackson, NSW

Distribution. Q, NSW, V, T, SA, WA: 0-100 m

***Cratena* Bergh, 1864**

Cratena Bergh, 1864: 198, 213

Type species. *Doris peregrina* Gmelin, 1791

***Cratena lineata* (Eliot, 1905)**

Facelina lineata Eliot, 1905: 288-289, pl. 16, figs. 4-5, pl. 17, figs 10-11

Cratena lineata.— Marshall & Willan, 1999: 255, fig. 266 (photo)

Cratena lineata.— Coleman, 2001: 110 (photo)

Type locality. Prison Island, Zanzibar, Africa

Distribution. Q, NSW, V, SA, WA, Indo-West Pacific: 0-18 m

***Echinopsole* Macnae, 1954**

Echinopsole Macnae, 1954: 25-26

Type species. *Echinopsole fulvus* Macnae, 1954

***Echinopsole breviceratae* Burn, 1962**

Echinopsole breviceratae Burn, 1962c: 124-125, figs 25-26

Echinopsole breviceratae.— Burn, 1989: pl. 57.4 (photo)

Type locality. Torquay, Victoria

Distribution. V, SA: 0-10 m

***Facelina* Alder & Hancock, 1855**

Facelina Alder & Hancock, 1855: XXII

Type species. *Eolida coronata* Forbes & Goodsir, 1839 = *Doris auriculata* Müller, 1776

***Facelina hartleyi* Burn, 1962**

Facelina hartleyi Burn, 1962c: 116, fig 17

Austraeolis fucia Burn, 1962c: 122-3, figs. 23-24

Phidiana hartleyi.— Coleman, 2001: 111 (photo)

Type locality. Flinders, Victoria

Distribution. V, T, SA: 0-12 m

***Facelina newcombi* (Angas, 1864)**

Flabellina newcombi Angas, 1864: 68-69, pl. 6, fig. 8

Phidiana newcombi.— Coleman, 2001: 112 (photo)

Type locality. Coodgee Bay, Sydney, NSW

Distribution. NSW, V: 0-10 m

Facelina sp 1

Distribution. V: 0 m

White body with yellow spots on sides and dorsum, blue line on long tail, cerata with yellowish stripes over brown digestive gland.

Facelina sp 2

Distribution. V: 0 m

Species with pink body becoming deep mauve at extremities, cerata white over red digestive gland, rhinophores roughened, foot corners very wide.

Facelina sp 3

Distribution. V: 0 m

Similar to *Facelina newcombi* in size and to *F. hartleyi* in colouration, but easily separated from both by sparsely papillate (not annulate) rhinophores and white outlining of cerata.

Favorinus J.E.Gray in M.E.Gray, 1850

Favorinus J.E.Gray in M. E. Gray, 1850b: 109

Type species. *Eolis alba* Alder & Hancock, 1844 (non *Eolidia alba* van Hasselt, 1824) = *Doris branchialis* Rathke, 1806

Favorinus pannuceus Burn, 1962

Favorinus pannuceus Burn, 1962c: 117, fig 18

Type locality. Flinders, Victoria

Distribution. V: 0-10 m

Palisa Edmunds, 1964

Palisa Edmunds, 1964:12

Type species. *Palisa papillata* Edmunds, 1964

Palisa sp

Palisa sp.— Coleman, 2001: 111 (photo: 'Ghostly Palisa')

Distribution. V: 0 m

Phyllodesmium Ehrenberg, 1831

Phyllodesmium Ehrenberg, 1831: signature h₃

Type species. *Phyllodesmium hyalinum* Ehrenberg, 1831

Phyllodesmium macphersonae (Burn, 1962)

Cratena macphersonae Burn, 1962c: 118-119, text figs. 19, 20

Phyllodesmium macphersonae.— Burn, 1989: pl. 56.6 (photo)

Phyllodesmium macphersonae.— Marshall & Willan, 1999: 257, fig. 274 (photo)

Phyllodesmium macphersonae.— Coleman, 2001: 113 (photo)

Type locality. Flinders pier, Victoria

Distribution. Q, NSW, V, T, SA, WA, Indo-West Pacific: 0-13 m

Phyllodesmium poindimiei (Risbec, 1928)

Aeolidia poindimiei Risbec, 1928: 246-247, text fig. 87, pl.9, fig 3

Phyllodesmium poindimiei.— Wells & Bryce, 1993: 160, 161

(photo: species 209)

Phyllodesmium poindimiei.— Marshall & Willan, 1999: 257, fig. 275 (photo)

Phyllodesmium poindimiei.— Coleman, 2001: 113 (photo)

Type locality. Poidimié, New Caledonia

Distribution. Q, NSW, V, SA, WA, Indo-West Pacific: 0-25 m

Phyllodesmium serratum (Baba, 1949)

Hervia serrata Baba, 1949: 105, 179, pl. 46, figs. 156-157, text figs. 142, 143

Phyllodesmium serratum.— Burn, 1989: pl. 57.1 (photo).— Wells & Bryce, 1993: 157,158 (photo species 205).— Coleman, 2001: 113-4 (photo)

Type locality. Sagami Bay, Japan

Distribution. Q, NSW, V, T, SA, WA, NT, Indo-West Pacific: 0-20 m

Family Embletoniidae Pruvot-Fol, 1954**Embletonia Alder & Hancock, 1851**

Embletonia Alder & Hancock, 1851: fam. 3, genus 14, pl.38

Type species. *Pterochilus pulcher* Alder & Hancock, 1844

Wägele & Willan (2000:170) refer to *Embletonia* as a "problematic" genus needing further study to resolve its relationships and systematic position. Miller (1977) and Miller & Willan (1991) transferred *Embletonia* and its family to the suborder Dendronotina. Rudman (1998:1015) retained them in the suborder Aeolidina, which is provisionally followed here.

Embletonia gracilis Risbec, 1928

Embletonia gracile Risbec, 1928: 271

Embletonia gracilis.— Miller & Willan, 1991: 2

Type locality. Baie de Canala, New Caledonia

Distribution. Q, NSW, V, T, Indo-West Pacific: 0-127 m

The species name was originally written with the neuter termination *-e*. However as *Embletonia* is feminine in gender, the appropriate termination is *-is*. A review of the literature concerning this name suggests that more than one species may be involved: a small (7-10 mm long) species with few pairs of cerata, to which Victorian specimens belong, and an elongate (to 40 mm) species with many pairs of cerata. Both types of animals are illustrated in Miller & Willan (1991).

Family Tergipedidae Bergh, 1889**Tergipes Cuvier, 1805**

Tergipes Cuvier, 1805: 433

Type species. *Limax tergipes* Forsskål, 1775

Tergipes sp

Distribution. V, SA, WA: 0-5 m

Possibly to be identified with *Tergipes tergipes* (Forsskål, 1775) of Atlantic distribution. Larger specimens (>7 mm) consistently possess two cerata in the right and left anterior liver branches, whereas this is unusual in European specimens.

Trinchesia von Ihering, 1879

Trinchesia Ihering, 1879: 137

Type species. *Doris caerulea* Montagu, 1804

Use of *Trinchesia* in place of *Cuthona* Alder & Hancock, 1855 follows Miller (2004).

Trinchesia anulata (Baba, 1949)

Cratena anulata Baba, 1949: 98-9, 175, pl.41 fig 145, textfigs 126, 127

Cuthona anulata.— Coleman, 2001: 107 (photo)

Type locality. Sagami Bay, Japan, 0-16 m

Distribution. V, Indo-West Pacific: 0-20 m

***Trinchesia catachroma* (Burn, 1963)**

Catriona catachroma Burn, 1963b: 15-16, figs 1-6
Cuthona catachroma.— Coleman, 2001: 107-8 (photo)

Type locality. Point Danger, Torquay, Victoria

Distribution. V: 0-12 m

***Trinchesia ornata* (Baba, 1937)**

Cuthona (Hervia) ornata Baba, 1937: 331, pl.2, fig.4
Cuthona ornata.— Marshall & Willan, 1999: 251, fig.254

Type locality. Tomioka, Amakusa, Japan

Distribution. Q, V, Japan, southern Africa: 0-10 m

***Trinchesia sororum* Burn, 1964**

Trinchesia sororum Burn, 1964b: 17-18, figs 11-15

Type locality. Point Lonsdale, Victoria

Distribution. V: 0 m

***Trinchesia thelmae* (Burn, 1964)**

Toorna thelmae Burn, 1964b: 20-21, figs 16-21
Cuthona thelmae.— Coleman, 2001: 108 (photo)

Type locality. Point Lonsdale, Victoria

Distribution. V, SA: 0-10 m

***Trinchesia viridiana* (Burn, 1962)**

Catriona viridiana Burn, 1962c: 111-112, text fig 13
Cuthona viridiana.— Coleman, 2001: 108 (photo)

Type locality. Point Danger, Torquay, Victoria

Distribution. V: 0-10 m

***Trinchesia* sp 1**

Distribution. V: 0 m

White body ornamented with blood-red spots and patches on sides and dorsum.

***Trinchesia* sp 2**

Cuthona sp.— Coleman, 2001: 108 (photo: 'Blue-headed Cuthona')

Distribution. V: 0-12 m

Cerata yellow with a bright blue subapical spot. Figured in Coleman (2001:108) as "Blue-headed Cuthona".

***Trinchesia* sp 3**

Cuthona sp.— Coleman, 2001: 108 (photo: 'Seagrass Cuthona')

Distribution. V, T, SA: 0-10 m

A flattish fawn and orange species adapted to life between the leaves of the seagrass *Amphibolis antarctica*.

***Trinchesia* sp 4**

Distribution. V: 0 m

Known only from preserved material. Cerata black.

***Trinchesia* sp 5**

Distribution. V: 0-2 m

Very small species (to 7 mm), colourless or palest yellowish, digestive gland fawn to brown, speckled with dark brown dots, strongly visible in body and cerata. At times, common in association with the hydroids *Obelia dichotoma* and *Monothecca*

flexuosa upon which it lives, eats and lays a small 1½ - 2 coiled sausage of shining white eggs.

***Trinchesia* sp 6**

Distribution. V: 15 m

Known from preserved specimens only. Very similar to *Trinchesia zelandica* (Odhner, 1924) from the Auckland Islands, south of New Zealand.

***Trinchesia* sp 7**

Distribution. V: 0 m

Cerata white with subapical blue band, below which a yellow spot.

***Trinchesia* sp 8**

Distribution. V, SA: 0 m

Cerata white with bright white net-like pattern all over, body with pale blue diamond-shaped patches along dorsum.

***Trinchesia* sp 9**

Distribution. V: 0 m

Cerata with bright pink digestive gland and ochre yellow tip.

***Trinchesia* sp 10**

Distribution. V: 0-10 m

Smaller and broader than *Trinchesia catachroma*, but with similar colour pattern, cerata inflated basally, with four rings of yellow.

***Trinchesia* sp 11**

Distribution. V: 0-12 m

Differentiated from preceding species by the wiggly or spirally twisted rhinophores, and the cerata with white tip below which a yellow and a blue band.

***Trinchesia* sp 12**

Distribution. V: 0-12 m

A very slender dull white species to 8 mm long, with long rhinophores, relatively few small, yellow and brown cerata, and a long tail, all overlaid with small silvery-white spots.

***Trinchesia* sp 13**

Distribution. V: 0-20 m

Body pale pink or orange, cerata with blue patch at mid-length and dark red apex, rhinophores irregularly shaped.

***Trinchesia* sp 14**

Distribution. V: 0-10 m

Body mauve, tips of rhinophores, tentacles and tail reddish-purple, basal third of cerata opaque white above which a well-separated yellow and a cream ring, foot corners rounded. *Facelina* sp. 2 is similarly, but more strongly, coloured, and is readily distinguished by very wide tentaculiform foot-corners.

***Trinchesia* sp 15**

Distribution. V: 5-8 m

Head and body with reddish-brown maculations, rhinophores with yellow band below clear tip, cerata long, crooked like an elbow, cream becoming pinkish sub-apically.

Family Fionidae Gray, 1857

Fiona Alder & Hancock in Forbes & Hanley, 1851

Fiona Alder & Hancock in Forbes & Hanley, 1851: contents, x, note

Type species. *Oithona nobilis* Alder & Hancock in Forbes & Hanley, 1851 = *Eolidia pinnata* Eschscholtz, 1831

Fiona pinnata (Eschscholtz, 1831)

Eolidia alba van Hasselt, 1824: 23

Eolidia pinnata Eschscholtz, 1831: 14, pl.19, fig 1

Fiona pinnata.—Coleman, 2001: 116 (photo)

Type locality. Sitka, Alaska

Distribution. Pelagic, world-wide, washed ashore after storms: 0 m

Acknowledgements

This checklist is dedicated to my late wife Margaret (20.6.1937-29.8.2004) whose patience and forbearance in no small way made it possible for me to actively pursue an obsession with opisthobranchs throughout family life and business activities.

Two people deserve more than special thanks. J. Hope Macpherson (Mrs Ian Black), then Conchologist, National Museum of Victoria, offered guidance, encouragement, and access to collections and library to an adolescent who evinced an interest in the then greatly neglected molluscan group, the Opisthobranchia. She still keeps an eye on all things malacological. Her successor, the late Dr Brian J. Smith, Curator of Molluscs, was equally helpful, securing type specimens and images of Australian species held in overseas institutions, as well as pertinent literature not available in Australia.

The writer is deeply indebted to the Museum Victoria and its many curatorial and staff members who have greatly helped with access to collection and library, to the many members of the Victorian Branch of the Malacological Society of Australasia, and of the Marine Research Group of Victoria (now part of the Field Naturalists Club of Victoria) for specimens collected during field-work, and to diving friends who aided by collecting specimens far below the intertidal collector. Each made it possible to compile this checklist.

Special thanks go the Robin Wilson and Liz Dane for facilitating the production of the checklist, to Clarrie Handreck for organising much of the field-work, to Neville Coleman, Jan Watson, Dave Staples, Ian Kirwen, Paul Dalton, Glenys Greenwood, John Chuk, Andrew Newton, Joan Hales, Leon Altoff, Audrey Falconer, and Platon Vafiadis for collecting and photographing many of the species, and to David Howlett for assiduously working the San Remo area and discovering many still unnamed species. Richard Willan kindly read an earlier version of the manuscript, and greatly improved both content and presentation. Finally, despite suffering time in the field in search of opisthobranchs, and proof-reading a copy of the manuscript, friend and colleague Ken Bell offered much advice on all kinds of molluscan and literature problems.

Bibliography

This bibliography comprises (a) the primary reference for every genus and species entry in the checklist, and (b) an as complete as possible list of other papers, articles and books that in some way cover species of the opisthobranch fauna of Victoria and the Bass Strait area.

- Abildgaard, P.C. 1791. Nyer efterretning om det skaldyr fra middelhavet, som Forskål har beskrevet under noun of Amonia tridentata. *Skrivter af Naturhistorie-Selskabet* 1(2): 171-175.
- Abraham, P. S. 1876. Notes on some genera of nudibranchiate Mollusca, with notices of a new genus and some hitherto undescribed species in the collection of the British Museum. *Annals and Magazine of Natural History* 18: 132-146.
- Abraham, P. S. 1877. Revision of the anthobranchiate nudibranchiate Mollusca, with descriptions or notices of forty-one hitherto undescribed species. *Proceedings of the Zoological Society of London*: 196-269.
- Adams, A. 1850. Monograph of the family Bullidae. In Sowerby, G.B.2 II., *Thesaurus Conchyliorum, or figures and descriptions of Recent shells*. 2(11): 553-608.
- Adams, A. 1855. Monographs of *Actaeon* and *Solidula*, two genera of gastropodous Mollusca, from the Cummingian collection. *Proceedings of the Zoological Society of London* 22: 58-62.
- Adams, H. & A. 1853-1858. *The genera of Recent Mollusca, arranged according to their organization*. Vol. 2. Van Voorst: London. 1: VI-XL, 1-484, 482: 481-661, 483: pls. 481-138 pp.
- Adams, A., & Angas, G. F. 1864. Descriptions of new species of shells, chiefly from Australia, in the Collection of Mr. Angas. *Proceedings of the Zoological Society of London* 1864: 35-40.
- Alder, J., & Hancock, A. 1843. Notice of a British species of *Calliopaea*, D'Orbigny, and of four new species of *Eolis*, with observations on the development and structure of the nudibranchiate Mollusca. *Annals and Magazine of Natural History* 12: 233-238.
- Alder, J., & Hancock, A. 1845-1855. *A Monograph of the British nudibranchiate Mollusca. Parts 1 -7*. Ray Society: London.
- Alder, J., & Hancock, A. 1854. Notice of some new species of British Nudibranchiata. *Annals and Magazine of Natural History* (2) 14: 102-105.
- Alder, J., & Hancock, A. 1862. Descriptions of a new genus and some new species of naked Mollusca. *Annals and Magazine of Natural History* (3) 10: 261-265.
- Alder, J., & Hancock, A. 1864. Notice of a collection of nudibranchiate mollusca made in India by Walter Elliot, Esq., with descriptions of several new genera and species. *Transactions of the Zoological Society of London* 5(3): 113-147.
- Allan, J. K. 1932. Australian nudibranchs. *Australian Zoologist* 7(2): 87-105.
- Allan, J. K. 1933. Opisthobranchs from Australia. *Records of the Australian Museum* 18(9): 443-450.
- Allan, J. K. 1934. Egg-cases of sea-snails and sea-slugs. *Victorian Naturalist* 50: 229-235.
- Allan, J. K. 1947. Nudibranchia from the Clarence River Heads, north coast, New South Wales. *Records of the Australian Museum* 21(8): 433-463.
- Angas, G. F. 1864. Description d'espèces nouvelles appartenant à plusieurs genres de Mollusques nudibranches des environs de Port Jackson, (Nouvelles-Galles du Sud), accompagnée de dessins faits d'après nature. *Journal de Conchyliologie* 12: 43-70.
- Angas, G. F. 1865. On the marine molluscan fauna of the Province of South Australia, with a list of all the species known up to the present time, together with remarks on their habitats and distribution, etc. *Proceedings of the Zoological Society of London* 1865: 155-190.
- Angas, G. F. 1867. List of species of marine Mollusca found in Port Jackson Harbour, New South Wales, and on the adjacent coasts, with notes on their habits, etc. Part 1. *Proceedings of the Zoological Society of London* 1867: 185-233.
- Angas, G. F. 1871a. Descriptions of thirty-four new species of shells from Australia. *Proceedings of the Zoological Society of London* 1871: 13-21.
- Angas, G. F. 1871b. A list of additional species of marine Mollusca to be included in the fauna of Port Jackson and adjacent coasts of New

- South Wales. *Proceedings of the Zoological Society of London* 1871: 87-101.
- Angas, G. F. 1877. Descriptions of one genus and twenty five species of marine shells from New South Wales. *Proceedings of the Zoological Society of London*: 171-177.
- Ascanius. 1772. *Philine quadripartita*, ein soust unbekanntes Seethier abgezeichnet und beschriebener. *Kongliga Svenska Vetenskapsakademiens Handlingar* 33: 329-331.
- Baba, K. 1930. Studies on Japanese nudibranchs. 3. *Venus* 2(3): 117-125.
- Baba, K. 1937. Opisthobranchia of Japan 2. *Journal of the Department of Agriculture, Kyushu Imperial University* 5(7): 289-344.
- Baba, K. 1938. Opisthobranchia of Kii, Middle Japan. *Journal of the Department of Agriculture, Kyushu Imperial University* 6(1): 1-19.
- Baba, K. 1949. *Opisthobranchia of Sagami Bay collected by His Majesty the Emperor of Japan*. Iwanami Shoten: Tokyo. 194 pp.
- Baba, K. 1955. *Opisthobranchia of Sagami Bay Supplement*. Iwanami Shoten: Tokyo. 59 pp.
- Baba, K. 1960. The genera *Okenia*, *Goniodoridella* and *Goniodoris* from Japan (Nudibranchia-Goniodorididae). *Publications of the Seto Marine Biological Laboratory* 8: 79-83.
- Baba, K. 1961a. On the identification and the affinity of *Tamanovalva limax*, a bivalved sacoglossan mollusc in Japan. *Publications of the Seto Marine Biological Laboratory* 9(1): 37-62.
- Baba, K. 1961b. The shells and radulae in *Berthelinia*, a bivalved sacoglossan genus. *Venus* 21(4): 389-401.
- Basedow, H., & Hedley, C. 1905. South Australian nudibranchs, and an enumeration of the known Australian species. *Transactions of the Royal Society of South Australia* 29: 134-160.
- Bebbington, A. 1977. Aplysiid species from Eastern Australia with notes on the Pacific Ocean Aplysiomorpha (Gastropoda: Opisthobranchia). *Transactions of the Zoological Society of London* 34: 87-147.
- Beddome, R.H. 1883. Description of some new marine shells of Tasmania. *Papers and Proceedings of the Royal Society of Tasmania* 1882: 167-170.
- Beesley, P. L., Ross, G. J. B., & Wells, A. (eds) 1998. *Mollusca: The Southern Synthesis. Fauna of Australia*. Vol. 5. CSIRO Publishing: Melbourne.
- Behrens, D. W. 2004. Pacific Coast Nudibranchs, Supplement 11. New species to the Pacific Coast and new information on the oldies. *Proceedings of the California Academy of Sciences* 55(2): 11-54.
- Bergh, L.S.R. 1860. Om Forekomsten af Neldfiim hos Mollusker. *Videnskabelige Meddeleser fra Dansk Naturhistorisk Forening*: 309-331.
- Bergh, L.S.R. 1864. Anatomiske Bidrag til Kundskab om aeolidierne. *Det Kongelige Danske Videnskaberne Selskabs Skrifter. Naturvidenskabelig og Mathematisk Afdeling*. (5) 7: 139-316.
- Bergh, L.S.R. 1867. *Phidiana lynceus* og *Ismaila monstrosa*. *Videnskabelige Meddeleser fra Dansk Naturhistorisk Forening* No. 7-9: 97-130.
- Bergh, L.S.R. 1869. Bidrag til en Monographi af Phyllidierne. *Naturhistorisk Tidsskrift Stiftet af Henrik Kroyer. Kjøbenhavn* (3) 5: 357-542.
- Bergh, L.S.R. 1873. Neue Nacktschnecken der Südsee, Malacologische Untersuchungen. *Journal des Museum Godeffroy* 1(2): 65-96 (137-168).
- Bergh, L.S.R. 1876. Malacologische Untersuchungen. In Semper, C.G. (ed.). *Reisen im Archipel der Philippinen von Dr. C. Semper* 2. 2(10): 377-427.
- Bergh, L.S.R. 1877a. Kritische Untersuchung der Ehrenberg'schen Doriden. *Jahrbücher der Deutschen Malakozoologischen Gesellschaft* 4(1): 45-76.
- Bergh, L.S.R. 1877b. Malacologische Untersuchungen. In Semper, C.G. (ed.). *Reisen im Archipel der Philippinen von Dr. C. Semper* 2. 2(12): 495-546.
- Bergh, L.S.R. 1877c. Malacologische Untersuchungen. In Semper, C.G. (ed.). *Reisen im Archipel der Philippinen von Dr. C. Semper* 2. 2(11): 429-494.
- Bergh, L.S.R. 1878. Malacologische Untersuchungen. In Semper, C.G. (ed.). *Reisen im Archipel der Philippinen von Dr. C. Semper* 2. 2(14): 603-645.
- Bergh, L.S.R. 1879a. Beiträge zu einer Monographie der Polyceraden. 1. *Verhandlungen der königlich-kaiserlich Zoologisch-botanischen Gesellschaft in Wien* 29: 599-632.
- Bergh, L.S.R. 1879b. Gattungen Nordische Doriden. *Archiv für Naturgeschichte* 45(1): 340-369.
- Bergh, L.S.R. 1879c. Neue Chromodoriden. *Malakozoologische Blätter Neue Folge* 1: 87-116.
- Bergh, L.S.R. 1880a. Malacologische Untersuchungen. In Semper, C.G. (ed.). *Reisen im Archipel der Philippinen von Dr. C. Semper* 2. 4(1): Suppl. (1) 1-78.
- Bergh, L.S.R. 1880b. Die Doriopsen des Mittelmeeres. *Jahrbuch der Deutschen Malakozoologischen Gesellschaft* 7: 297-328.
- Bergh, L.S.R. 1884a. Report on the Nudibranchiata. *Challenger Reports. Zoology* 10(26): 1-154.
- Bergh, L.S.R. 1884b. Malacologische Untersuchungen. In Semper, C.G. (ed.). *Reisen im Archipel der Philippinen von Dr. C. Semper* 2. 3(15): 647-754.
- Bergh, L.S.R. 1888. Malacologische Untersuchungen. In Semper, C.G. (ed.). *Reisen im Archipel der Philippinen von Dr. C. Semper* 2. 3(16): 755-814.
- Bergh, L.S.R. 1900. Ergebniss einer Reise nach dem Pacific (Schauinsland 1896-1897) Die Opisthobranchier. *Zoologische Jahrbücher* 13(3): 207-246.
- Bergh, L.S.R. 1904. Malacologische Untersuchungen. In Semper, C.G. (ed.). *Reisen im Archipel der Philippinen von Dr. C. Semper* 9. 6(1): 1-56.
- Bergh, L.S.R. 1905. Malacologische Untersuchungen. In Semper, C.G. (ed.). *Reisen im Archipel der Philippinen von Dr. C. Semper* 9. 6(2): 57-115.
- Bertsch, H. 1977. The Chromodoridinae nudibranchs from the Pacific coast of America. Part 1. Investigative methods and supraspecific taxonomy. *Veliger* 20: 107-118.
- Beu, A. G. 2004. Marine Mollusca of oxygen isotope stages of the last 2 million years in New Zealand. Part 1: Revised generic positions and recognition of warm-water and cool-water migrants. *Journal of the Royal Society of New Zealand* 34(2): 111-265.
- Blainville, M.H. de. 1816. Quatrième mémoire sur les mollusques, de l'ordre des cyclobranches. *Bulletin des Sciences par la Société Philomathique*: 93-97.
- Blainville, M.H. de. 1817. *Dictionnaire des Sciences Naturelles Supplement 5*.
- Blainville, M.H. de. 1821. Mémoire sur le genre Hyale. *Journal de Physique, de chimie, et d'histoire naturelle élémentaires* 93: 81-97.
- Blainville, M.H. de. 1824. *Dictionnaire des Sciences Naturelles*. Vol. 32. Levrault: Paris & Strasbourg. 567 pp.
- Blainville, M. H. de. 1825-1827. *Manuel de Malacologie et de Conchyliologie*. Pp. 1-647 (1827), 649-664 (1827), pl. 1-87 (1827).
- Boas, J. E. V. 1886. Spolia Atlantica. Bidrag til Pteropodernes. Morphologi og Systematik am til Kundskaben om deres Geografiske Udbredelse. *Det Kongelige Danske Videnskaberne Selskabs Skifter, 6 Raekke, Naturvidenskabelig og Mathematisk Afdeling* 4: 1-248.
- Bosc, L.A.G. 1817. *Nouveau dictionnaire d'histoire naturelle*. Vol. 7. Deterville: Paris. 586 pp.
- Bouchet, P., & Rocroi, J.-P. 2005. Classification and nomenclature of gastropod families. *Malacologia* 47(1-2): 1-397.
- Boyd, S. 1999. The introduced Mollusca of Port Phillip Bay. Pp. 129-149 in: Hewitt, C.L., Campbell, M.L., Thresher, R.E. & Martin, R.B. (eds), *Marine Biological Invasions of Port Phillip Bay, Victoria*. Centre for Research on Introduced Marine Pests. Technical Report No. 20. CSIRO Marine Research: Hobart. 344pp.
- Brodie, G. D. 2005. Redescription of the Australian endemic nudibranch *Dendrodoris maugeana* Burn, 1962 (Gastropoda: Opisthobranchia: Dorididae): new and reviewed features important for future phylogenetic analyses of porostomes. *Molluscan Research* 25(1): 37-46.
- Brodie, G. D., & Calado, G. 2006. *Dendrodoris arborescens* (Collingwood, 1881) (Mollusca: Nudibranchia): larval characteristics reveal a masked porostome species. *Records of the Western Australian Museum*, Supplement 69: 119-126.
- Brodie, G.D., Willan, R. C., & Collins, J.D. 1997. Taxonomy and occurrence of *Dendrodoris nigra* and *Dendrodoris fumata* (Nudibranchia: Dendrodorididae) in the Indo-west Pacific region. *Journal of Molluscan Studies* 63: 407-423.

- Brown, T. 1827. *Illustrations of the Conchology of Great Britain and Ireland*. Edinburgh. Preface + 53 pls.
- Burn, R. 1957a. A new species of Opisthobranchia from Victoria (Mollusca: Gastropoda). *Victorian Naturalist* 74: 115-117.
- Burn, R. 1957b. On some Opisthobranchia from Victoria. *Journal of the Malacological Society of Australia* 1(1): 11-29.
- Burn, R. 1958. Further Victorian Opisthobranchia. *Journal of the Malacological Society of Australia* 1(2): 20-36.
- Burn, R. 1959. Comments on the Australian umbraculacean Mollusca. *Journal of the Malacological Society of Australia* 1(3): 28-30.
- Burn, R. 1960a. A bivalve gastropod. *Nature* 186: 179.
- Burn, R. 1960b. Australian bivalve gastropods. *Nature* 187: 44-46.
- Burn, R. 1960c. New names for two Victorian opisthobranch molluscs. *Journal of the Malacological Society of Australia* 1(4): 70.
- Burn, R. 1961a. A common Victorian nudibranch [*Ceratosoma brevicaudatum*]. *Victorian Naturalist* 77(11): 316-317.
- Burn, R. 1961b. *Drepaniella mapae* gen. et spec. nov., a new goniodorid nudibranch from south-eastern Australia. *Veliger* 3(4): 102-104.
- Burn, R. 1961c. A new dorid nudibranch from Torquay, Victoria. *Veliger* 4(2): 55-56.
- Burn, R. 1961d. *Eucrairia* nom. nov. for *Drepaniella* Burn. *Veliger* 4(1): 51.
- Burn, R. 1962a. Notes on a collection of Nudibranchia (Gastropoda: Dorididae and Dendrodorididae) from South Australia with remarks on the species of Basedow and Hedley, 1905. *Memoirs of the National Museum of Victoria* 25: 149-172, pl. 141.
- Burn, R. 1962b. On the new pleurobranch subfamily Berthellinae (Mollusca: Gastropoda); a revision and new classification of the species of New South Wales and Victoria. *Memoirs of the National Museum of Victoria* 25: 129-148.
- Burn, R. 1962c. Descriptions of Victorian nudibranchiate Mollusca, with a comprehensive review of the Eolidacea. *Memoirs of the National Museum of Victoria* 25: 95-128.
- Burn, R. 1963a. Australian Runcinacea (Mollusca: Gastropoda). *Australian Zoologist* 13: 9-22.
- Burn, R. 1963b. Descriptions of Australian Eolidacea (Mollusca: Opisthobranchia) 1. The genera *Catriona* and *Herviella*. *Journal of the Malacological Society of Australia* 1(7): 12-20.
- Burn, R. 1964a. Australian Sea Slugs. *Australian Natural History* 14(12): 396-399.
- Burn, R. 1964b. Descriptions of Australian Eolidacea (Mollusca: Opisthobranchia) 2. The genera *Nossis*, *Eubranthus*, *Trinchesia* and *Toorna*. *Journal of the Malacological Society of Australia* 1(8): 10-22.
- Burn, R. 1965a. A centennial commentary and zoogeographical remarks on Angas' Sydney nudibranchs (Mollusca: Gastropoda). *Journal de Conchyliologie* 104: 85-93.
- Burn, R. 1965b. Rediscovery and taxonomy of *Edentellina typica* Gatliff & Gabriel. *Nature* 206: 735-736.
- Burn, R. 1966a. On three new Chromodoridinae from Australia (Opisthobranchia: Nudibranchia). *Veliger* 8: 191-197.
- Burn, R. 1966b. The identity of Maplestone's opisthobranchs. *Victorian Naturalist* 83(4): 76-78.
- Burn, R. 1966c. Notes on some opisthobranchs mainly from South Australia. *Records of the South Australian Museum* 15(2): 329-352.
- Burn, R. 1966d. The opisthobranchs of a caulerpan microfauna from Fiji. *Proceedings of the Malacological Society of London* 37(1): 45-65.
- Burn, R. 1966e. Descriptions of Australian Eolidacea (Mollusca: Opisthobranchia) 3. The genera *Tularia*, *Embletonia* and *Austraeolis*, with a note on *Tergipes pauculus* Burn, 1962. *Journal of the Malacological Society of Australia* 1(9): 25-35.
- Burn, R. 1966f. Opisthobranchia. Port Phillip Survey 1957-1963. *Memoirs of the National Museum of Victoria* 27: 265-288.
- Burn, R. 1967a. Descriptions of two new species of *Okenia* (Nudibranchia: Doridacea) from south-eastern Australia. *Proceedings of the Royal Zoological Society of New South Wales* 1965-66: 52-58.
- Burn, R. 1967b. First record of a pelagic eolid from Victoria. *Victorian Naturalist* 84(4): 116-117.
- Burn, R. 1969. A memorial report on the Tom Crawford collection of Victorian Opisthobranchia. *Journal of the Malacological Society of Australia* 1(12): 64-106.
- Burn, R. 1970. Eight additional opisthobranch molluscs for New South Wales. *Proceedings of the Royal Zoological Society of New South Wales* 1968-69: 51-54.
- Burn, R. 1972a. The genus *Polycera* in Australian waters. *Australian Newsletter (Malacological Society of Australia) New Series* 16: 8-9.
- Burn, R. 1972b. A guide to the Ascoglossa or sap-sucking sea-slugs of Australia. *Australian Natural History* 17(5): 174-178.
- Burn, R. 1973a. Pearson Island Expedition 1969 - Opisthobranchs. *Transactions of the Royal Society of South Australia* 97(3): 201-205.
- Burn, R. 1973b. *Limapontia* in Australia. *Australian Shell News (Newsletter of the Malacological Society of Australia)* 4: 2.
- Burn, R. 1974a. Notes on some benthonic opisthobranchs from Port Phillip Bay, Victoria. *Journal of the Malacological Society of Australia* 3(1): 43-57.
- Burn, R. 1974b. *Limapontia* in New Zealand waters. *Australian Shell News* 7: 5.
- Burn, R. 1975a. Records of three opisthobranchs from South Australia. *Journal of the Malacological Society of Australia* 3(2): 62.
- Burn, R. 1975b. Distributional data for two south-eastern Australian opisthobranchs. *Journal of the Malacological Society of Australia* 3(2): 88.
- Burn, R. 1975c. Notes on *Paliolla cooki* (Angas, 1864) from southern Australia. *Journal of the Malacological Society of Australia* 3(2): 107-110.
- Burn, R. 1975d. A list of dorid nudibranchs of Australia. Pp.514-517, in Thompson, T.E. 1975 Dorid nudibranchs from eastern Australia. *Journal of Zoology, London* 176: 477-517.
- Burn, R. 1977. Tropical opisthobranchs in South Australia. *Australian Shell News No.* 19: 8.
- Burn, R. 1978a. *Marianina rosea* (Provot-Fol, 1930): further records from Australia. *Journal of the Malacological Society of Australia* 4(1-2): 28.
- Burn, R. 1978b. A review of Australian species of *Austrocylichna*, *Nipponatys*, *Cylichnatys* and *Diniatys* (Mollusca: Gastropoda: Haminoeidae). *Journal of the Malacological Society of Australia* 4: 93-112.
- Burn, R. 1978c. Records of *Philineopsis lineolata* (H. & A. Adams, 1854) from south-eastern Australia. *Journal of the Malacological Society of Australia* 4(1-2): 6.
- Burn, R. 1979. New locality records for *Lophopleurella wilsoni* (Tate, 1889) from southern Australia. *Journal of the Malacological Society of Australia* 4(3): 128.
- Burn, R. 1984. Opisthobranchia. P. 67-74 in: Phillips, D.A.B., Handreck, C., Bock, P.E., Burn, R., Smith, B.J. & Staples, D.A. (eds), *Coastal Invertebrates of Victoria: an atlas of selected species*. Marine Research Group of Victoria/Museum of Victoria: Melbourne.
- Burn, R. 1989. Opisthobranchs (Subclass Opisthobranchia). Pp. 725-788 in: Shepherd, S.A. & Thomas, I.M. (eds), *Marine Invertebrates of Southern Australia. Part II*. South Australian Government Printing Division: Adelaide.
- Burn, R. 1990. An annotated list of opisthobranch molluscs from San Remo, Westernport, Victoria. *Marine News (Newsletter of the Marine Research Group of Victoria)* 100: 9-15.
- Burn, R. 1998. Orders Acochlidea, Rhodopemomorpha, Sacoglossa. Pp. 959-974 in: Beesley, P.L., Ross, G.J.B. & Wells, A. (eds), *Mollusca: the Southern Synthesis. Fauna of Australia Volume 5*. CSIRO Publishing: Melbourne.
- Burn, R., & Bell, K. N. 1974a. Description of *Retusa chrysome* Burn sp nov. (Opisthobranchia) and its food resources from Corner Inlet, Victoria. *Memoirs of the National Museum of Victoria* 35: 115-119.
- Burn, R., & Bell, K. N. 1974b. Description of *Retusa pelyx* Burn sp nov. and its food resources from Swan Bay, Victoria. *Journal of the Malacological Society of Australia* 3: 37-42.
- Burn, R., & Bell, K.N. 1976. Cryptic molluscs inhabiting *Galeolaria* in Victoria. *Victorian Naturalist* 93(6): 232-236.
- Burn, R., & Miller, M. C. 1969. A new genus, *Caldukia*, and an extended description of the type species, *Proctonotus ? affinis* Burn, 1958 (Mollusca: Gastropoda: Arminacea: Antiopellidae). *Journal of the Malacological Society of Australia* 1(12): 23-31.
- Cantraine, F. J. 1835. Mollusques - "diagnoses ou descriptions succintes de quelques espèces nouvelles de mollusques." *Bulletin de Academie Royale de Sciences, Bruxelles* 2(10): 380-401.

- Cheeseman, T.F. 1881. On some new species of nudibranchiate Mollusca. *Transactions and Proceedings of the New Zealand Institute* 13: 222-224.
- Cheeseman, T.F. 1886. On a new species of *Chromodoris*. *Transactions and Proceedings of the New Zealand Institute* 18: 137.
- Coleman, N. 1975. *What Shell is That?* Paul Hamlyn: Sydney. 308 pp.
- Coleman, N. 1976. *Shell Collecting in Australia*. A.H. & A.W. Reed: Sydney. 176 pp.
- Coleman, N. 1977. *A Field Guide to Australian Marine Life*. Rigby Limited: Adelaide. 223 pp.
- Coleman, N. 1981a. *A Field Guide to the Marine Life of South-Eastern Australia*. Rigby Publishers Limited: Adelaide. 167 pp.
- Coleman, N. 1981b. *Shells Alive!* Rigby Publishers Limited: Adelaide. 96 pp.
- Coleman, N. 1989. *Nudibranchs of the South Pacific*. Vol. 1. Sea Australia Resource Centre: Springwood, Queensland. 64 pp.
- Coleman, N. 2001. *1001 Nudibranchs - Catalogue of Indo-Pacific Sea Slugs*. Neville Coleman's Underwater Geographic Pty. Ltd.: Springwood, Queensland.
- Collingwood, C. 1881. On some new species of nudibranchiate mollusca from the eastern seas. *Transactions of the Linnean Society of London, Zoology* 2: 123-140.
- Cotton, B. C., & Godfrey, F. K. 1933. South Australian shells (Including descriptions of new genus and species). Part 7. *South Australian Naturalist* 14(3): 72-108.
- Crosse, H., & Fischer, P. 1865. Description d'espèces nouvelles de l'Australie méridionale. *Journal de Conchyliologie* 13: 38-55.
- Cuvier, G. 1797. Sur un nouveau genre de mollusque. *Bulletin des Sciences de la Société Philomathique de Paris* 1(1): 105.
- Cuvier, G. 1804. Mémoire sur la Phyllidie et sur le Pleurobranche, deux nouveaux genres de mollusques de l'ordre des gastéropodes, et voisins des patelles et des oscabriens, dont l'un est nu et dont l'autre porte une coquille cachée. *Annales du Muséum d'Histoire Naturelle* 5(28): 266-276.
- Cuvier, G. 1805. Mémoire sur la *Scyllée*, l'*Eolide* et le *Glaucus* avec des additions au mémoire sur la *Tritonie*. *Annales du Muséum d'Histoire Naturelle* 6: 416-436.
- Cuvier, G. 1817. *Le Règne Animal*. Vol. 2.
- Dall. 1871. Descriptions of sixty new forms of mollusks from the west coast of North America and the North Pacific Ocean, with notes on others already described. *American Journal of Conchology* 7(2): 93-160.
- Dall. 1902. Illustrations and descriptions of new, unfigured, or imperfectly known shells, chiefly American, in the U.S. National Museum. *Proceedings of the United States National Museum* 24: 449-566.
- Day, J. H., & Hutchings, P. A. 1984. Descriptive notes on the fauna and flora of Merimbula, Pambula and Back Lakes, New South Wales. *Australian Zoologist* 21(3): 269-289.
- Dayrat, B. 2005. Advantages of naming species under the *PhyloCode*: an example of how a new species of *Discodorididae* (Mollusca, Gastropoda, *Euthyneura*, *Nudibranchia*, *Doridina*) may be named. *Marine Biology Research* 1: 216-232.
- Dayrat, B. 2006. A taxonomic revision of *Paradoris* sea slugs (Mollusca, Gastropoda, *Nudibranchia*, *Doridina*). *Zoological Journal of the Linnean Society* 147: 125-238.
- Deshayes, G.P. 1835-1845. *Histoire naturelle des animaux sans vertèbres*. Deuxième édition. Vol. 1-11.
- Deshayes, G.P. 1839-1858. *Traité élémentaire de Conchyliologie, avec les applications de cette science à la Géologie. Atlas (1839-1853): Explication des Planches: 1-80*.
- Dorgan, K. M., Valdés, A., & Gosliner, T. M. 2002. Phylogenetic systematics of the genus *Platydoris* (Mollusca, *Nudibranchia*, *Doridoidea*) with descriptions of six new species. *Zoologica Scripta* 31: 271-319.
- Eales, N. B. 1960. Revision of the world species of *Aplysia* (Gastropoda, *Opisthobranchia*). *Bulletin of the British Museum (Natural History)*. *Zoology* 5(10): 268-404.
- Edgar, G. J. 1997. *Australian Marine Life, the plants and animals of temperate waters*. Reed Books: Kew, Australia. 544 pp.
- Edmunds, M. 1964. Eolid Mollusca from Jamaica, with descriptions of two new genera and three new species. *Bulletin of Marine Science of the Gulf and Caribbean* 14(1): 1-32.
- Ehrenberg, C.G. 1828-1831. *Symbolae physicae seu icones et descriptiones animalium evertebratorum sepositis insectis quae ex itinere per Africam Borealem et Asiam Occidentalem - novae aut illustratae redierunt. Decas I Mollusca*. Pages unnumbered.
- Eliot, C. N. E. 1902. On some nudibranchs from Zanzibar. *Proceedings of the Zoological Society (London)*: 62-72.
- Eliot, C. N. E. 1904. On some nudibranchs from East Africa and Zanzibar. Part 3. *Proceedings of the Zoological Society (London)*: 354-385.
- Eliot, C. N. E. 1905. On some nudibranchs from East Africa and Zanzibar. Part 6. *Proceedings of the Zoological Society (London)*: 268-298.
- Eliot, C. N. E. 1909. Notes on a collection of nudibranchs from Ceylon. *Spoila Zeylanica* 6(23): 79-95.
- Eschscholtz, J. F. von. 1831. *Zoologischer Atlas - Beschreibungen neuer Thierarten - Zweiter Reise um die Welt*. Part 4. 19 pp.
- Fahey, S. J., & Gosliner, T. M. 2003. Mistaken identities: on the *Discodorididae* genera *Hoplordoris* Bergh, 1880 and *Carminodoris* Bergh, 1889 (*Opisthobranchia*, *Nudibranchia*). *Proceedings of the California Academy of Sciences* 54(10): 169-208.
- Fahey, S. J., & Gosliner, T. M. 2004. A phylogenetic analysis of the *Aegiridae* Fischer, 1883 (Mollusca, *Nudibranchia*, *Phanerobranchia*) with descriptions of eight new species and a reassessment of phanerobranch relationships. *Proceedings of the California Academy of Sciences* 55(34): 613-589.
- Fahey, S. J., & Valdés, A. 2005. Review of *Acanthodoris* Gray, 1850 with a phylogenetic analysis of *Onchidorididae* Alder & Hancock, 1845 (Mollusca, *Nudibranchia*). *Proceedings of the California Academy of Sciences* 56(20): 213-273.
- Férussac, A. E. J. d'Audebard de. 1822. *Tableaux systématiques des animaux mollusques suivis d'un Prodrome général pour tous les mollusques terrestres ou fluviaux vivants ou fossiles*. Premier partie, *Tableaux systématiques généraux*. Anthus-Bertrand: Paris. i- xlvii pp.
- Fleming, J. 1823. On a reversed species of *Fusus* (*Fusus retroversus*). *Memoirs of the Wernerian Natural History Society* 4(2): 498-500.
- Fleming, J. 1828. *A History of British animals, exhibiting the descriptive characters and systematical arrangement of the genera and species of quadrupeds, birds, reptiles, fishes, Mollusca and Radiata of the United Kingdom, etc.*: Edinburgh.
- Forbes, E. 1838. *Malacologia monensis, a catalogue of the Mollusca inhabiting the Isle of Man and the neighboring sea*. 63 pp.
- Forbes, E. 1844. Report on the mollusca and radiata of the Aegean Sea, and on their distribution, considered as bearing on geology. *Report of the 13th Meeting of the British Association for the Advancement of Science (1843)*: 131-139.
- Forbes, E., & Goodsir, J. 1839. Notice of zoological researches in Orkney and Shetland during the month of June, 1839. *The Athenaeum, Journal of Literature, Science and Fine Arts* No. 618: 647.
- Forbes, E., & Hanley, S. 1851. *A History of British Mollusca, and Their Shells*. Vol. 3. 562-612 pp.
- Forster, J.G.A. 1777. A voyage around the world in H.M.S. Resolution, commanded by Capt. J. Cook, during 1772-5. 1: 49.
- Freame, M.E. 1935. A sea-slug (*Dorid*). *Victorian Naturalist* 51: 244.
- Gabb, W. M. 1873. Description of some new genera of Mollusca. *Proceedings of the Academy of Natural Sciences of Philadelphia* 1872: 270-274.
- Gabriel, C. J. 1933. The Victorian "Umbrella Shell", *Umbraculum corticale* Tate. *Victorian Naturalist* 50(2): 48.
- Gabriel, C. J. 1936. *Victorian Sea Shells*. FNCV Handbook: Melbourne. 68 pp.
- Gabriel, C. J. 1962. Additions to the marine molluscan fauna of south eastern Australia including descriptions of new genus *Pillarginella*, six new marine species and two subspecies. *Memoirs of the National Museum of Victoria* 25: 177-210.
- Gardiner, A. P. 1936. Engel's paper on "The English species of the family Pleurobranchidae". *Journal of Conchology* 20(7): 195-198.
- Gascoigne, T. 1974. A note of some sacoglossan penial styles (Gastropoda: *Opisthobranchia*). *Zoological Journal of the Linnean Society* 55(1): 53-59.
- Gascoigne, T. 1976. The reproductive systems and classification of the *Stiligeridae* (*Opisthobranchia*: *Sacoglossa*). *Journal of the Malacological Society of Australia* 3(3-4): 157-172.

- Gascoigne, T., & Sartory, P. K. 1974. The teeth of three bivalved gastropods and some other species of the order Sacoglossa. *Proceedings of the Malacological Society of London* 41: 109-124.
- Gatliff, J. H. 1888. A list of some of the shells of the marine Mollusca found upon the Victorian coast. Part II. *Victorian Naturalist* 5: 111-114.
- Gatliff, J. H., & Gabriel, C. J. 1908a. On some new species of Victorian marine Mollusca. *Proceedings of the Royal Society of Victoria* 21(1): 365-367.
- Gatliff, J. H., & Gabriel, C. J. 1908b. Additions to and revision of the Catalogue of Victorian marine Mollusca. *Proceedings of the Royal Society of Victoria* 21(1): 368-391.
- Gatliff, J. H., & Gabriel, C. J. 1909. Additions to the Catalogue of the marine shells of Victoria. *Proceedings of the Royal Society of Victoria* 22(1): 37-46.
- Gatliff, J. H., & Gabriel, C. J. 1911. On some new species of Victorian marine mollusca. *Proceedings of the Royal Society of Victoria* 24(1): 187-192.
- Gatliff, J. H., & Gabriel, C. J. 1913a. On some new species and varieties of Victorian marine Mollusca. *Proceedings of the Royal Society of Victoria* 26(1): 67-70.
- Gatliff, J. H., & Gabriel, C. J. 1913b. Additions to the Catalogue of marine shells of Victoria. *Proceedings of the Royal Society of Victoria* 26(1): 71-87.
- Gatliff, J. H., & Gabriel, C. J. 1917. Additions to and alterations in the Catalogue of the marine shells of Victoria. *Proceedings of the Royal Society of Victoria* 30(1): 21-31.
- Gatliff, J. H., & Gabriel, C. J. 1922. Additions to and alterations in the Catalogue of Victorian marine Mollusca. *Proceedings of the Royal Society of Victoria* 34(2): 128-161.
- Gillett, K., & Yaldwyn, J. C. 1969. *Australian Seashores in Colour*. A.H. & A.W. Reed: Sydney. 112 pp.
- Gistel, J. von N.F.X. 1848. *Naturgeschichte des Thierreichs für höhere Schulen*.
- Gmelin, J.F. 1791. *Caroli a Linné, Systema naturae per regna tria naturae. Editio decima tertia*. Leipzig, Germany: 1(6) class 6, Vermes: 3021-3910.
- Goddard, J. H. R. 2005. Ametamorphic direct development in *Dendrodoxia behrensi* (Nudibranchia: Dendrodoxiidae), with a review of developmental mode in the family. *Proceedings of the California Academy of Sciences* 56(19): 201-211.
- Gosliner, T. M. 1987. Review of the Nudibranch genus *Melibe* (Opisthobranchia: Dendronotacea) with descriptions of two new species. *The Veliger* 29: 400-414.
- Gosliner, T. M. 1989. Revision of the Gastropteridae (Opisthobranchia: Cephalaspidea) with descriptions of a new genus and six new species. *The Veliger* 32: 333-381.
- Gosliner, T. M. 1995. Introduction and spread of *Philine auriformis* (Gastropoda: Opisthobranchia) from New Zealand to San Francisco Bay and Bodega Harbor. *Marine Biology* 122: 249-255.
- Gosliner, T. M. 2006. Marine Gastropoda collected by the Steamer Albatros from the Philippines in 1908. *Records of the Western Australian Museum*, Supplement 69: 83-93.
- Gosliner, T. M., & Johnson, S. 1994. Review of the genus *Hallaxa* (Nudibranchia: Actinocyclusidae) with description of nine new species. *The Veliger* 37(2): 155-191.
- Gosliner, T. M., & Smith, V. G. 2003. Systematic review and phylogenetic analysis of the nudibranch genus *Melibe* (Opisthobranchia: Dendronotacea) with descriptions of three new species. *Proceedings of the California Academy of Sciences* 130: 83-181.
- Gould, A. A. 1859. Descriptions of shells collected in the North Pacific Exploring Expedition under Captains Ringgold and Rodgers. *Proceedings of the Boston Society of Natural History* 7: 138-142.
- Grande, C., Templado, J., Cervera, J. L., & Zardoya, R. 2004. Phylogenetic relationships among Opisthobranchia (Mollusca: Gastropoda) based on mitochondrial *cox1*, *trnV*, and *rnl* genes. *Molecular Phylogenetics and Evolution* 33: 378-388.
- Gray, J.E. 1825. A list and description of some species of shells not taken notice of by Lamarck. *Annals of Philosophy* 25: 134-140, 407-415.
- Gray, J.E. 1843. Shells. In Dieffenbach, E. *Travels in New Zealand, with contributions to the geography, geology, botany and natural history of that country*. 2, 396pp. Murray: London.
- Gray, J.E. 1847. A list of the genera of recent Mollusca, their synonymia and types. *Proceedings of the Zoological Society of London* 15: 129-219.
- Gray, J.E. 1850a. *Catalogue of the Mollusca in the Collection of the British Museum II. II. Pteropoda*: 1-45. E. Newman: London.
- Gray, M. E. 1850b. *Figures of molluscous animals, selected from various authors*. Vol. 4. 124 pp.
- Griffith, E., & Pidgeon, E. 1833. The Mollusca and Radiata. In Griffith, E. *The Animal Kingdom by Cuvier* Part 38: 1-192.
- Hasselt, J. C. van. 1824. Uittreksel uit eenen brief van Dr J.C. Hasselt, aan Prof. van Swinderen. *Algemene Konst-en Letterbode 1824* 2: 20-24; 3: 34-39; 4: 54-55.
- Haszprunar, G., & Hess, A. 2005. A new *Rhodope* from the Roscoff area (Bretagne), with a review of *Rhodope* species (Gastropoda: Nudibranchia?). *Spixiana* 28(3): 193-197.
- Hedley, C. 1894. On some naked Australian Marine Mollusca Part I. *Proceedings of the Linnean Society of New South Wales* (2) 9: 126-128.
- Hedley, C. 1895. On a molluscan genus new to, and another forgotten from Australia. *Proceedings of the Royal Society of Victoria (n.s.)* 7: 197-200.
- Hedley, C. 1902a. Studies on Australian Mollusca. Part V. *Proceedings of the Linnean Society of New South Wales* 26: 700-708.
- Hedley, C. 1902b. Studies on the Australian Mollusca. Part VI. *Proceedings of the Linnean Society of New South Wales* 26(1): 7-29.
- Hedley, C. 1903. Mollusca. Part 2. Scaphopoda and Gastropoda. Scientific results of the trawling expedition of H.M.C.S. Thetis, off the coast of New South Wales, in February and March, 1898. *Australian Museum Memoir* 4: 327-402.
- Hedley, C. 1912. Descriptions of some new or noteworthy shells in the Australian Museum. *Records of the Australian Museum* 8(3): 131-160.
- Hedley, C. 1918. A check-list of the marine fauna of New South Wales. Part 1. Mollusca. *Journal and Proceedings of the Royal Society of New South Wales* 51: M1-M120.
- Hedley, C. 1920. Concerning *Edentellina*. *Proceedings of the Malacological Society of London* 14: 74-76.
- Hedley, C. 1923. Studies on Australian Mollusca. Part 14. *Proceedings of the Linnean Society of New South Wales* 48(3): 301-316.
- Hedley, C., & May, W.L. 1908. Mollusca from one hundred fathoms seven miles east of Cape Pillar, Tasmania. *Records of the Australian Museum* 7: 108-125.
- Hewitt, C. L., Campbell, M. L., Thresher, R. E., & Martin, R. B. 1999. *Marine Biological Invasions of Port Phillip Bay, Victoria*. Centre for Research on Introduced Marine Pests. Technical Report No. 20. CSIRO Marine Research: Hobart. 344 pp.
- Hinds. 1844. Descriptions of new species of *Ringicula* and *Neaera*, from the cabinets of Sir E. Belcher and Hugh Cuming Esq. *Proceedings of the Zoological Society of London* 12: 96-98.
- Ihering, H. von. 1879. Einiges neue über Mollusken. *Zoologischer Anzeiger* 2: 136-138.
- International Commission on Zoological Nomenclature. 1999. *International Code of Zoological Nomenclature. Fourth Edition*. The International Trust for Zoological Nomenclature: London. 306 pp.
- International Commission on Zoological Nomenclature. 2000. Opinion 1942. *Haminoea* [Turton] in Turton & Kingston in Carrington, 1830 and *Haminoeinae* Pilsbry, 1895 (Mollusca, Gastropoda): placed on Official Lists as correct spellings. *Bulletin of Zoological Nomenclature* 57(1): 52-53.
- Iredale, T. 1925. Mollusca from the continental shelf of eastern Australia. *Records of the Australian Museum* 14(4): 243-270.
- Iredale, T. 1929. Strange molluscs in Sydney Harbour. *Australian Zoologist* 5(4): 337-352.
- Iredale, T. 1930. Some notable name changes. *Australian Zoologist* 6: 175.
- Iredale, T. 1936. Australian molluscan notes. No. 2. *Records of the Australian Museum* 19: 267-340.
- Iredale, T., & O'Donoghue, C. H. 1923. List of the British nudibranchiate Mollusca. *Proceedings of the Malacological Society of London* 15: 195-233.
- Iredale, T., & McMichael, D.F. 1962. A reference list of the marine Mollusca of New South Wales. *Australian Museum, Sydney: Memoir XI*: 1-109.

- Jansen, P. 2000. *Seashells of South-East Australia*. Capricornica Publications: Lindfield. 117 pp.
- Jeffreys, J.G. 1877. New and peculiar Mollusca of the Eulimidae and other families of Gastropoda, as well as of the Pteropoda, procured in the "Valorous" expedition. *Annals and Magazine of Natural History* (4) 19: 317-339.
- Jensen, K. R. 1985. Annotated checklist of Hong Kong Ascoglossa (Mollusca: Opisthobranchia) with descriptions of four new species. P. 77-107 in: Morton, J.E. & Dudgeon, D. (eds), *The malacofauna of Hong Kong and southern China 2. Proceedings of the Second International Workshop on the Malacofauna of Hong Kong and Southern China, Hong Kong 6-24 April 1983*. Hong Kong University Press: Hong Kong.
- Jensen, K. R. 1993. Sacoglossa (Mollusca, Opisthobranchia) from Rottneest Island and central Western Australia. Pp. 207-253 in: Wells, F.E., Walker, D.I., Kirkman, H. & Lethbridge, R. (eds), *Proceedings of the Fifth International Marine Biological Workshop: The Marine Flora and Fauna of Rottneest Island, Western Australia. 1*. Western Australian Museum: Perth.
- Jensen, K. R. 1997. *Sacoglossa systematik, fylogeni og evolution* (Mollusca, Opisthobranchia). Vestjydsk Forlag: Copenhagen.
- Jensen, K. R., & Wells, F. E. 1990. Sacoglossa (=Ascoglossa) (Mollusca, Opisthobranchia) from southern Western Australia. Pp. 297-331 in: Wells, F.E., Walker, D.I., Kirkman, H. & Lethbridge, R. (eds), *Proceedings of the Third International Marine Biological Workshop: The Marine Flora and Fauna of Albany, Western Australia. 1*. Western Australian Museum: Perth.
- Kawaguti, S., & Baba, K. 1959. A preliminary note on a two-valved sacoglossan gastropod, *Tamanovalva limax*, n. gen., n. sp., from Tamano, Japan. *Biological Journal of Okayama University* 5: 177-184.
- Kay, E. A. 1968. A review of the bivalved gastropods and a discussion of evolution within the Sacoglossa. *Symposia of the Zoological Society of London* 22: 109-134.
- Kay, E. A., & Young, D. K. 1969. The Doridacea (Opisthobranchia; Mollusca) of the Hawaiian Islands. *Pacific Science* 23: 172-231.
- Keferstein, W. 1862. Kopt-Weichthiere: Malacozoa Cephalota. In Bronn's. *Klassen und Ordnungen des Thierreichs* 3(2): 523-1500.
- Kershaw, R.C. 1953. A systematic list of the Mollusca of Tasmania, Australia. *Papers and Proceedings of the Royal Society of Tasmania* 89: 289-355.
- Klussman-Kolb, A. 2004. Phylogeny of the Aplysiidae (Gastropoda, Opisthobranchia) with new aspects of the evolution of seahares. *Zoologica Scripta* 33: 439-462.
- Klussman-Kolb, A., & Dinapoli, A. 2006. Systematic position of the pelagic Thecosomata and Gymnosomata within Opisthobranchia (Mollusca, Gastropoda) - a revival of the Pteropoda. *Journal of Zoological Systematics & Evolutionary Research* 44(2): 118-129.
- Koelliker, A. 1847. *Rhodope*, nuovo genere di Gasteropodi. *Giornale dell' Imperiale Reale Istituto Lombardo de Scienze, Lettere ed Arti* (e *Biblioteca Italiana*). 16(47): 239-249.
- Kosse, J.F.J. 1813. *Dissertatio de Pteropodum ordine et novo ipsius genera*. Halle. Pp. 10-16.
- Kuroda, T., & Habe, T. 1952. *Check list and bibliography of the Recent Mollusca of Japan*. Leo W. Stach: Tokyo. 210 pp.
- Lemche, H. 1967. *Rhinodiaphana g.n. ventricosa* (Jeffreys, 1865) redescribed (Gastropoda Tectibranchiata). *Sarsia* 29: 207-214.
- Leue, S. 1813. De *Pleurobranchaea* novo molluscorum genere. *Dissrn. Inaug. Acad. Halle 1813*: 1-13.
- Lightfoot, J. 1786. *A Catalogue of the Portland Museum, lately the property of the Duchess Dowager of Portland, deceased, which will be sold at auction, by Mr Skinner and Co.*: London. viii & 194 pp.
- Linnaeus, C. 1758. *Systema Naturae per regna tria naturae. Editio decima, reformata*.
- Linnaeus, C. 1767. *Systema Naturae per regna tria naturae. Editio duodecima, reformata*.
- Lovén, S.L. 1844. Om nordiska hafs - mollusker. *Ofversigt af Kongl. Vetenskaps Akademiens Förhandlingar. Stockholm* 1(3): 48-53.
- Lovén, S.L. 1846. *Index molluscorum litora Scandinaviae occidentalia habitantium*. 1-28 pp.
- Macnae, W. 1954. On some Eolidacean Nudibranchiate Molluscs from South Africa. *Annals of the Natal Museum* 13(1): 1-50.
- Macpherson, J. H., & Chapple, E.H. 1951. A systematic list of the marine and estuarine Mollusca of Victoria. *Memoirs of the National Museum of Victoria* 17: 107-156.
- Macpherson, J. H., & Gabriel, C. J. 1962. *Marine Molluscs of Victoria*. Melbourne University Press & National Museum of Victoria: Melbourne xv. 475 pp.
- Marcus, Ev. du B.-R. 1972. On some opisthobranchs from Florida. *Bulletin of Marine Science* 22: 284-308.
- Marcus, Ev. du B.-R. 1982. Systematics of the genera of the order Ascoglossa (Gastropoda). *Journal of Molluscan Studies, Supplement* 10: 1- 31.
- Marcus, Ev. & Er. 1963. Opisthobranchs from the Lesser Antilles. *Studies on the Fauna of Curaçao and other Carribean Islands* 19(79): 1-76.
- Marcus, Er. 1955. Opisthobranchia from Brazil. *Boletim da Faculdade de Filosofia Ciências e Letras da Universidade de São Paulo, Zoologia* 20: 89-261.
- Marcus, Er. 1964. A new species of *Polycera* (Nudibranchia) from California. *Nautilus* 77: 128-131.
- Marshall, J.G., & Willan, R. C. 1999. *Nudibranchs of Heron Island, Great Barrier Reef. A Survey of the Opisthobranchia (Sea Slugs) of Heron and Wistari Reefs*. Backhuys: Leiden.
- Martens, E. von. 1879. Übersicht der von ihm (W. Peters) von 1843 bis 1847 in Mossambique gessammelten Mollusca. *Monatsberichte der Königlichen.-Preussische Akademie der Wissenschaften zu Berlin* 31: 727-749.
- May, W.L. 1903. On Tenison-Woods types in the Tasmanian Museum, Hobart. *Proceedings of the Royal Society of Tasmania, 1902*: 106-114.
- May, W.L. 1921. *A Check-list of the Mollusca of Tasmania*. Government Printer: Tasmania. 114 pp.
- May, W.L. 1923a. *An Illustrated Index of Tasmanian Shells: with 47 plates and 1052 species*. Government Printer: Hobart. 100 pp.
- May, W.L. 1923b. Mollusca of King Island, with descriptions of five new species. *Papers and Proceedings of the Royal Society of Tasmania, 1923*: 47-55.
- McCoy, F. 1885. Natural History of Victoria. Plate 104. *Cetorhinus maximus* (Linn. sp.) The Basking Shark. *Prodromus of the Zoology of Victoria* 2(Decade XI): 11-15.
- Medina, M., & Walsh, P. J. 2000. Molecular systematics of the order Anaspidea based on mitochondrial DNA sequence (12S, 16S and C01). *Molecular Phylogenetics and Evolution* 15(1): 41-58.
- Meisenheimer, J. 1903. Über eine neue Familie der gymnosomen Pteropoden aus dem Material der Deutschen Tiefsee Expedition (Pterocaniden). *Zoologischer Anzeiger* 26: 92-99.
- Meisenheimer, J. 1905. Pteropoda. *Wissenschaftliche Ergebnisse der Deutschen Tiefsee-Expedition auf dem Dampfer "Valdivia" 1898-1899*. 9(1): 1-314.
- Meisenheimer, J. 1906. Die Pteropoden der deutschen Süd-polar Expedition 1901-1903. *Deutsche Südpolar-Expedition 1901-1903. 9(Zoologie)* 1(2): 92-152.
- Menke, K. T. 1830. *Synopsis methodica molluscorum generum omnium et specierum earum, quae in Museo Menkeano adservantur etc.* Pyrmonti. 169 pp.
- Mikkelsen, P. M. 1996. The evolutionary relationships of Cephalaspidea S. I. (Gastropoda: Opisthobranchia): a phylogenetic analysis. *Malacologia* 37: 375-442.
- Mikkelsen, P. M. 2002. Shelled opisthobranchs. *Advances in Marine Biology* 42: 67-136.
- Millen, S.V., & Martynov, A. 2005. Redescriptions of the nudibranch genera *Akiodoris* Bergh, 1879 and *Armodoris* Minichev, 1972 (Suborder Doridacea), with a new species of *Akiodoris* and a new family Akiodorididae. *Proceedings of the California Academy of Sciences* 56(1): 1-22.
- Miller, M. C. 1971. Aeolid nudibranchs (Gastropoda: Opisthobranchia) of the family Flabellinidae and Eubranchidae from New Zealand waters. *Zoological Journal of the Linnean Society* 50: 311-337.
- Miller, M. C. 1977. Aeolid nudibranchs (Gastropoda: Opisthobranchia) of the family Tergipedidae from New Zealand waters. *Zoological Journal of the Linnean Society* 60: 197-222.
- Miller, M. C. 1988. *Aeolidia helicochorda*, a new aeolid nudibranch (Gastropoda: Opisthobranchia) from New Zealand. *New Zealand Journal of Zoology* 14: 391-397.

- Miller, M. C. 1991. On the identity of the dorid nudibranch *Homoiodoris novaezelandiae* Bergh, 1904 (Gastropoda: Opisthobranchia). *Journal of Natural History* 25: 293-304.
- Miller, M. C. 1995. New species of the dorid nudibranch genus *Paradoris* Bergh, 1884 (Gastropoda: Opisthobranchia) from New Zealand. *Journal of Natural History* 29: 901-908.
- Miller, M. C. 1996. A new species of the dorid nudibranch genus *Polycera* Cuvier, 1816 (Gastropoda: Nudibranchia) from New Zealand. *Journal of Molluscan Studies* 62: 443-450.
- Miller, M. C. 2001. Aeolid nudibranchs (Gastropoda: Opisthobranchia) of the family Aeoliidae from New Zealand waters. *Journal of Natural History* 35: 629-662.
- Miller, M. C. 2004. An appraisal of the identity of New Zealand species of the aeolid nudibranch family Tergipedidae (Gastropoda: Opisthobranchia). *Journal of Natural History* 38: 1183-1192.
- Miller, M. C. 2005. A new species of the dorid nudibranch genus *Polycera* Cuvier, 1817 (Gastropoda, Opisthobranchia) from New Zealand. *Vita Malacologica* 3: 51-54.
- Miller, M. C., & Willan, R. C. 1986. A review of the New Zealand arminacean nudibranchs (Opisthobranchia: Arminacea). *New Zealand Journal of Zoology* 13: 377-408.
- Miller, M. C., & Willan, R. C. 1991. Redescription of *Embletonia gracile* Risbec, 1928 (Nudibranchia: Embletoniidae): relocation to suborder Dendronotacea with taxonomic and phylogenetic implications. *Journal of Molluscan Studies* 58: 1-12.
- Montagu, G. 1815. Descriptions of several new or rare animals, principally marine, discovered on the South Coast of Devonshire. *Transactions of the Linnean Society of London, Zoology* 11: 1-26.
- Montfort, D. de. 1810. *Conchyliologie Systematique, et classification methodique des coquilles*. Vol. 2. F. Schoell: Paris. 676 pp.
- Mörch, O.A.L. 1863. Contributions à la faune malacologique des Antilles Danoises. *Journal de Conchyliologie* (3) 11: 21-43.
- Müller, O.F. 1776. *Zoologia Danicae. Prodrum seu animalium Daniae et Nowegiae ingenarum characteres, nomina, et synonyma imprimis popularium*. I-XXXII, 282 pp.
- Nakano, R. 2004. *Opisthobranchs of the Japan Islands*. Rutles Inc.: Tokyo.
- Newman, L. J. 1998. Orders Thecosomata and Gymnosomata. Pp. 565-1234 in: Beesley, P.L., Ross, G.J.B. & Wells, A. (eds), *Mollusca: The Southern Synthesis. Fauna of Australia. Volume 5*. CSIRO: Melbourne.
- Newton, R.B. 1891. *Systematic list of the F.E. Edwards Collection of British Oligocene & Eocene Mollusca in the British Museum (Natural History)*. British Museum (Natural History): London. xxviii + 365 pp.
- Odhner, N. Hj. 1924. New Zealand Mollusca (Papers from Dr Th. Mortensen's Pacific Expedition 1914-16 XIX). *Videnskabelige Meddelelser fra Dansk Naturhistorisk Forening* 77: 1-90.
- Odhner, N. Hj. 1934. The Nudibranchia. *British Antarctic ("Terra Nova") Expedition 1910. Zoology* 7(5): 229-310.
- O'Donoghue, C. H. 1929. Report on the Opisthobranchiata. [in] Zoological results of the Cambridge Expedition to the Suez Canal, 1924. *Transactions of the Zoological Society of London* 22(6): 713-841.
- Oken, L. 1813-1826. *Lehrbuch der Naturgeschichte*. Vol. 3 (1). C.H.Reclam: Leipzig.
- Orbigny, A. d'. 1836. Voyage dans l'Amerique meridionale exécuté pendant les années 1826-1833. *Mollusques* 5(3): 49-184.
- Pease, W. H. 1860a. Descriptions of new species of Mollusca from the Sandwich Islands. *Proceedings of the Zoological Society of London* 28: 18-36.
- Pease, W. H. 1860b. Descriptions of new species of Mollusca from the Sandwich Islands. (Part 2). *Proceedings of the Zoological Society of London* 28: 141-148.
- Pease, W. H. 1861. Descriptions of new species of Mollusca from the Pacific Islands. *Proceedings of the Zoological Society of London for 1861*: 242-247.
- Pelseneer, P. 1886. Description d'un nouveau genre de ptéropode gymnosome (Notobranchaea). *Bulletin Scientifique (Historique et Littéraire) du Département du Nord. Lille. (2)* 9: 79-80.
- Pelseneer, P. 1887. Report on the Pteropoda collected by H.M.S. Challenger during the years 1873-76. Part I. The Gymnosomata. *Report of the Scientific Results of the Voyage of H.M.S. Challenger 1873-76, Zoology* 19: 1-74.
- Pelseneer, P. 1888. Report on the Pteropoda. II. The Thecosomata. *Challenger Reports. Zoology* 23: 1-132.
- Phipps, C. J. 1774. *A voyage towards the North Pole undertaken by his Majesty's Command 1773*. W. Bowyer & J. Nichols: London. 275 pp.
- Pilsbry, H. A. 1895. Tectibranchiata. In Tyron, G.W. *Manual of Conchology* 15: 134-436.
- Pilsbry, H. A. 1895-1896. Philinidae, Gastropteridae, Aglajidae, Aplysiidae, Oxynoeidae, Runcinidae, Umbraculidae, Pleurobranchidae. in: Tyron, G.W. *Manual of Conchology*.
- Pilsbry, H. A. 1921. Marine mollusks of Hawaii, parts 14-15. *Proceedings of the Academy of Natural Sciences of Philadelphia* 72: 360-382.
- Poore, G.C.B, Wilson, R., Goman, M., & Lu, C.C. 1985. Museum of Victoria Bass Strait Survey, 1979-1984. A final report to MRAAC (Marine Research Allocations Advisory Committee) on Marine Sciences and Technologies Grant Number 80/141. Melbourne: Museum of Victoria Council.
- Powell, A. W. B. 1937. New species of Nudibranchiate Mollusca from Auckland waters. *Records of the Auckland Institute and Museum* 2(2): 119-124.
- Pritchard, G.B., & Gatliff, J.H. 1898. Catalogue of the marine shells of Victoria. Part I. *Proceedings of the Royal Society of Victoria* 10(2): 236-284.
- Pritchard, G.B., & Gatliff, J. H. 1902. Catalogue of the marine shells of Victoria. Part VI. *Proceedings of the Royal Society of Victoria* 15(2): 176-223.
- Pruvot-Fol, A. 1930. Diagnose provisoire (imcompètes) des espèces nouvelles et liste provisoire des mollusques nudibranches recueillis par Mme. A. Pruvot-Fol en nouvelle Cadédonie (Ile de Pins.). *Bulletin de Muséum National d'Histoire Naturelle* (2) 2(2): 229-233.
- Pruvot-Fol, A. 1931. Notes de systématique sur le opisthobranches. *Bulletin de Muséum National d'Histoire Naturelle* (2) 3(3): 309-310.
- Pruvot-Fol, A. 1932. Notes sur quelques gymnosomes de provenances diverses et diagnose d'un genre nouveau. *Archives de Zoologie expérimentale de générale Paris* 74: 507-529.
- Pruvot-Fol, A. 1933. Mission Robert Ph. Dollfus en Egypte. Opisthobranchiata. *Mémoires de l'Institut d'Egypte* 21: 89-159.
- Pruvot-Fol, A. 1942. Les Gymnosomes. I. *Dana Reports* 4(20): 1-54.
- Pruvot-Fol, A. 1951. Étude des Nudibranches de la Méditerranée. *Archives de Zoologie Expérimentale et Générale* 88(1): 1-79.
- Pruvot-Fol, A. 1953. Étude de quelques Opisthobranches de la côte atlantique du Maroc et du Sénégal. *Travaux de l'Institut Scientifique Chérifien* 5: 3-95.
- Quoy, J. R. C., & Gaimard, J. P. 1827. Observations zoologiques faites à bord de "l'Astrolabe" en Mai 1826, dans le détroit de Gibraltar: description des genres Biphore, Carinaire, Hyale, Flèche, Cleodore, Anatifé et Briarée. *Annales de Sciences Naturelles* 10: 225-239.
- Quoy, J. R. C., & Gaimard, J. P. 1832-1833. Voyages de découvertes de l'Astrolabe exécuté par ordre du Roi pendant les années 1826-1829 sous le commandement de M.J Dumont d'Urville. *Zoologie* 2: 1-686.
- Rafinesque. 1814a. *Specchio delle Scienze o Giornale Enciclopedico de Sicilia*. Vol. 2. Rafinesque: Palermo. 196 pp.
- Rafinesque. 1814b. *Précis des découvertes somiologiques ou zoologiques et botaniques*.: Palermo. 30 pp.
- Rang, P.C.S.A.L. 1825. Description d'un genre nouveau de la classe des ptéropodes, et de deux espèces de ptéropodes. *Annales de Sciences Naturelles* 5: 283-287.
- Rang, P.C.S.A.L. 1827. Description de deux genres nouveaux (*Cuvieria* et *Euribia*) appartenant à la classe de Ptéropodes. *Annales de Sciences Naturelles* 12: 320-329.
- Rang, P.C.S.A.L. 1828a. Notice sur quelques mollusques nouveaux appartenant au genre Cléodore, et établissement et monographie du sous-genre Créseis. *Annales de Sciences Naturelles* 13: 302-319.
- Rang, P.C.S.A.L. 1828b. Histoire naturelle des Aplysiens. Première famille de l'ordre des Tectibranches. Pp. 1-84 in: de Férussac, B. (ed.) *Histoire Naturelle, Générale et Particulière des Mollusques*.: Paris.
- Rang, P.C.S.A.L. 1829. *Manuel de l'histoire naturelle des mollusques et de leurs coquilles*. Paris.
- Richmond, M.H. 1990. *Tasmanian Sea Shells common to other Australian States. [Revised edition, 1997]*. Richmond Printers: Devonport. 80 pp.
- Richmond, M.H. 1992. *Tasmanian Sea Shells Volume 2*. Richmond Printers: Devonport. 111 pp.
- Risbec, J. 1928. Contribution à l'étude des nudibranches Néo-Calédoniens. *Faune des Colonies Françaises* 2(1): 1-328.

- Risso, A. 1818. Mémoire sur quelques Gastéropodes nouveaux, Nudibranches et Tectibranches observé dans la mer de Nice. *Journal de Physique* 87: 368-377.
- Röding, P.F. 1798. *Museum Boltenianum*. Vol. 2. VIII + 199 pp.
- Rudman, W. B. 1968. Three new species of the opisthobranch family Aglajidae from New Zealand. *Transactions of the Royal Society of New Zealand* 10(23): 211-216.
- Rudman, W. B. 1970. A revision of the genus *Philine* in New Zealand with descriptions of two new species (Gastropoda Opisthobranchia). *Journal of the Malacological Society of Australia* 2(1): 23-34.
- Rudman, W. B. 1972a. The anatomy of the opisthobranch genus *Hydatina* and the functioning of the mantle cavity and alimentary canal. *Zoological Journal of the Linnean Society* 51(2): 121-139.
- Rudman, W. B. 1972b. On *Melanochlamys* Cheeseman, 1881, a genus of the Aglajidae (Opisthobranchia, Gastropoda). *Pacific Science* 26(1): 50-62.
- Rudman, W. B. 1972c. The genus *Philine* (Opisthobranchia, Gastropoda). *Proceedings of the Malacological Society of London* 40: 171-187.
- Rudman, W. B. 1974. A comparison of *Chelidonura*, *Navanax* and *Aglaja* with the other genera of the Aglajidae (Opisthobranchia: Gastropoda). *Zoological Journal of the Linnean Society* 54(3): 185-212.
- Rudman, W. B. 1978. The dorid opisthobranch genera *Halgerda* Bergh and *Sclerodoris* Eliot from the Indo-West Pacific. *Zoological Journal of the Linnean Society* 62: 59-88.
- Rudman, W. B. 1982. The taxonomy and biology of further aeolidacean and arminacean nudibranch molluscs with symbiotic zooxanthellae. *Zoological Journal of the Linnean Society* 74: 147-196.
- Rudman, W. B. 1983. The Chromodorididae (Opisthobranchia: Mollusca) of the Indo-West Pacific: *Chromodoris splendida*, *C. aspersa* and *Hypselodoris placida* colour groups. *Zoological Journal of the Linnean Society* 78: 105-173.
- Rudman, W. B. 1984. The Chromodorididae (Opisthobranchia: Mollusca) of the Indo-West Pacific: a review of the genera. *Zoological Journal of the Linnean Society* 81: 115-273.
- Rudman, W. B. 1986. The Chromodorididae (Opisthobranchia: Mollusca) of the Indo-West Pacific: *Noumea flava* colour groups. *Zoological Journal of the Linnean Society* 88: 377-404.
- Rudman, W. B. 1987a. The Chromodorididae (Opisthobranchia: Mollusca) of the Indo-West Pacific: *Chromodoris epicuria*, *C. aureopurpurea*, *C. annulata*, *C. coi*, and *Risbecia tryoni* colour groups. *Zoological Journal of the Linnean Society* 90: 305-407.
- Rudman, W. B. 1987b. The genus *Trapania* (Nudibranchia: Goniodorididae) in the Indo-West Pacific. *Journal of Molluscan Studies* 53: 189-212.
- Rudman, W. B. 1988. The Chromodorididae (Opisthobranchia: Mollusca) of the Indo-West Pacific: the genus *Ceratosoma* J.E. Gray. *Zoological Journal of the Linnean Society* 93: 133-185.
- Rudman, W. B. 1990. The Chromodorididae (Opisthobranchia: Mollusca) of the Indo-West Pacific: further species of *Glossodoris*, *Thorunna* and the *Chromodoris aureomarginata* colour group. *Zoological Journal of the Linnean Society* 100: 263-326.
- Rudman, W. B. 1991a. Further studies on the taxonomy and biology of the octocoral-feeding genus *Phyllodesmium* Ehrenberg, 1831 (Nudibranchia: Aeolidioidea). *Journal of Molluscan Studies* 57: 167-203.
- Rudman, W. B. 1991b. Purpose in Pattern: The evolution of colour in chromodorid nudibranchs. *Journal of Molluscan Studies, Supplement, T.E. Thompson Memorial Issue* 57(4): 5-21.
- Rudman, W. B. 1998. Opisthobranchia: History of Discovery, and Order Nudibranchia. Pp. 919-921, 990-1017 in: Beesley, P.L., Ross, G.J.B. & Wells, A. (eds), *Mollusca: The Southern Synthesis. Fauna of Australia. Volume 5*. CSIRO: Melbourne.
- Rudman, W. B. 2004. Further species of the opisthobranch genus *Okenia* (Nudibranchia: Goniodorididae) from the Indo-West Pacific. *Zootaxa* 695: 1-70.
- Rudman, W. B. 2006. Australian Museum, Sydney, *Sea Slug Forum* <http://www.seaslugforum.net>
- Rudman, W. B., & Avern, G. J. 1989. The genus *Rostanga* Bergh, 1879 (Nudibranchia: Dorididae) in the Indo-West Pacific. *Zoological Journal of the Linnean Society* 96: 281-338.
- Rüppell, E., & Leuckart, F. S. 1828. *Atlas zu der Reise im nordlichen Afrika von Eduard Rüppell. Neue wirbellose thiere des Rothen Meeres*. Brönnner: Frankfurt. 50 pp.
- Sars, G. O. 1878. *Bidrag til Kundskaben om Norges Arktiske Fauna I. Mollusca regionis arcticae Norvegiae*. Christiania Universitets: Christiania. 466 pp.
- Sars, M. 1870. Bidrag til Kundskab om Christiania-fjordens Fauna 2. Pp. 49-114 in: *Christiania-fjordens Mollusker*
- Schiøtte, T. 1998. A taxonomic revision of the genus *Diaphana* Brown, 1827, including a discussion of the phylogeny and zoogeography of the genus (Mollusca: Opisthobranchia). *Steenstrupia* 24(1): 77-140.
- Schumacher, C.F. 1817. *Essai d'un Nouveau Système des Habitations des Vers Testacés*. Schultzy: Copenhague. 287 pp.
- Shepherd, S. A., & Thomas, I. M. (eds) 1989. *Marine Invertebrates of Southern Australia. Part 2*. South Australian Government Printing Division: Adelaide.
- Singleton, F.A. 1937. The McCoy Society Expedition to Lady Julia Percy Island. 14. Mollusca. *Proceedings of the Royal Society of Victoria* 49(2): 387-396.
- Smith, B. J., Black, J. H., & Shepherd, S. A. 1989. Molluscan egg masses and capsules. Pp. 841-891. in: Shepherd, S.A. & Thomas, I.M. (eds), *Marine Invertebrates of Southern Australia Part II*. South Australian Government Printing Division: Adelaide.
- Smith, E. A. 1872. Remarks on several species of Bullidae, with descriptions of some hitherto undescribed forms, and of a new species *Planaxis*. *Annals and Magazine of Natural History* (4) 9: 344-355.
- Smith, E. A. 1891. Descriptions of new species of shells from the 'Challenger' Expedition. *Proceedings of the Zoological Society of London* 1891: 436-445.
- Smith, V. G., & Gosliner, T. M. 2005. A new species of *Marionia* (Gastropoda: Nudibranchia) from the Caroline Islands. *Proceedings of the California Academy of Sciences* 56(6): 66-75.
- Sowerby, G.B. II. 1868. Monograph of the genus *Bulla*. In: Reeve, L. *Conchologia Iconica* 16: pls. 1-6.
- Sowerby, G.B. II. 1869. Monograph of the genus *Aplysia*. in: Reeve, L. (ed.) *Conchologia Iconica: or, illustrations of the shell of molluscous animals*.
- Sowerby, G.B. II. 1870. Descriptions of forty-eight new species of shells. *Proceedings of the Zoological Society of London* 1870: 249-259.
- Stimpson, W. 1855a. Descriptions of some new marine invertebrata. *Proceedings of the Academy of Natural Sciences of Philadelphia* 7: 385-395.
- Stimpson, W. 1855b. Descriptions of some of the new marine invertebrata from the Chinese and Japanese Seas. *Proceedings of the Academy of Natural Sciences of Philadelphia* 7(10): 374-384.
- Suter, H. 1909. Descriptions of new species and subspecies of New Zealand Mollusca, with notes on a few species. *Proceedings of the Malacological Society of London* 8: 253-265.
- Swennen, C., & Dekker, R. 1995. *Corambe batava* Kerbert, 1886 (Gastropoda: Opisthobranchia), an immigrant in the Netherlands, with a revision of the family Corambidae. *Journal of Molluscan Studies* 61: 107-107.
- Tate, R. 1879. Descriptions of the marine Gastropoda of South Australia. *Transactions of the Royal Society of South Australia* 2: 138-140.
- Tate, R. 1889. Descriptions of some new species of marine Mollusca from South Australia and Victoria. *Transactions and Proceedings and Report of the Royal Society of South Australia* 11: 60-66.
- Tate, R., & May, W.L. 1901. A revised census of the marine Mollusca of Tasmania. *Proceedings of the Linnean Society of New South Wales* 26(3): 344-471.
- Tenison Woods, J.E. 1876. Description of new Tasmanian shells. *Papers and Proceedings of the Royal Society of Tasmania* 1875: 134-162.
- Tenison Woods, J.E. 1878. Census, with brief descriptions of the marine shells of Tasmania and the adjacent islands. *Papers and Proceedings of the Royal Society of Tasmania* 1877: 26-57.
- Thiele, J. 1925. Gastropoda der Deutschen Tiefsee-Expedition. *Wissenschaftliche Ergebnisse der Deutschen Tiefsee-Expedition* 21: 158-268.
- Thompson, T. E. 1970a. Eastern Australian Pleurobranchomorpha. *Journal of Zoology, London* 160: 173-198.

- Thompson, T. E. 1970b. Eastern Australian Dendronotoidea. *Zoological Journal of the Linnean Society* 51(1): 63-77.
- Thompson, T. E. 1972. Chromodorid nudibranchs from eastern Australia. *Journal of Zoology, London* 166: 391-409.
- Thompson, T. E. 1973. Sacoglossan gastropod molluscs from eastern Australia. *Proceedings of the Malacological Society of London* 40: 239-251.
- Thompson, T. E. 1975. Dorid nudibranchs from eastern Australia. *Journal of the Zoological Society, London* 176: 477-517.
- Thompson, T. E., & Bennett, I. 1970. Observations on Australian Glaucidae. *Zoological Journal of the Linnean Society* 49(3): 187-197.
- Thompson, T. E., & Brown, G. H. 1974. *Atagema gibba* Pruvot-Fol, a doridacean nudibranch new to the British fauna. *Journal of Conchology* 28: 233-237.
- Tokioka, T., & Baba, K. 1964. Four new species and a new genus of the family Gastropteridae from Japan (Gastropoda: Opisthobranchia). *Publications of the Seto Marine Biological Laboratory* 12: 201-229.
- Trinchese, S. 1872. Un nuovo genere de la famiglia delle Eolididae. *Annali del Museo avico di Storia naturale de Genova* 2: 86-132.
- Trinchese, S. 1876. Anatomia della *Caliphylla mediterranea*, Costa. *Rendiconto delle Sessioni della Accademia delle Scienze dell'Istituto di Bologna, Anno 1875-1875*: 84-87.
- Troschel, F.H. 1854. Beiträge zur Kenntnis der Pteropoden. *Archiv für Naturgeschichte* 20(1): 196-241.
- Turton, W. 1830. Conchology, arranged on the amended system. Pp. 21 unnumbered (signature E₂-G₆) in: Turton, W. & Kingston, J.F., in Carrington, N.T. (ed.). in: *The Teignmouth, Dawlish and Torquay guide, part 2 (The natural history of the district; or, lists of the different species of animals, vegetables and minerals, and their respective localities, scientifically arranged)*
- Vafiadis, P. 1999. Intertidal sighting of *Stiliger smaragdinus* Baba, 1949 - an uncommon mollusc. *Victorian Naturalist* 116(4): 118.
- Valdés, A. 2001a. Deep-sea cryptobranch dorid nudibranchs (Mollusca, Opisthobranchia) from the tropical West Pacific, with descriptions of two new genera and eighteen new species. *Malacologia* 43(1-2): 237-311.
- Valdés, A. 2001b. On the publication date, authorship and type species of *Umbraculum* and *Tylodina* (Gastropoda: Opisthobranchia: Tylodinoidea). *Nautilus* 115: 29-34.
- Valdés, A. 2002. A phylogenetic analysis and systematic revision of the cryptobranch dorids (Mollusca, Nudibranchia, Anthobranchia). *Zoological Journal of the Linnean Society* 136: 535-636.
- Valdés, A., & Bouchet, P. 2005. Cephalaspidea, Thecosomata, Gymnosomata, Aplysiomorpha, Umbraculida, Acochlidacea, Sacoglossa, Cyliodrobulida, Nudipleura. Pp 258-263, 279-280. In: Bouchet, P. & Rocroi, J.-P. 2005. Classification and nomenclator of gastropod families. *Malacologia* 47(1-2): 1-397.
- Valdés, A., & Campillo, O.A. 2004. Systematics of pelagic aeolid nudibranchs of the family Glaucidae (Mollusca, Gastropoda). *Bulletin of Marine Sciences* 75(3): 381-389.
- Valdés, A., & Gosliner, T.M. 2001. Systematics and phylogeny of the caryphyllidia-bearing dorids (Mollusca, Nudibranchia), with descriptions of a new genus and four new species from Indo-Pacific deep waters. *Zoological Journal of the Linnean Society* 133: 103-198.
- van der Spoel, S. 1967. *Euthecosomata, a group with remarkable developmental stages (Gastropoda, Pteropoda)*. Journal van Noorduijn en Zoon. N.V.: Gorinchem. 375 pp.
- van der Spoel, S. 1976. *Pseudotheosomata, Gymnosomata and Heteropoda (Gastropoda)*. Bohn, Scheltema, and Holkema: Utrecht. 484 pp. pp.
- van der Spoel, S. 1987. *Diacavolinia* nov. gen. separated from *Cavolinia* (Pteropoda, Gastropoda). *Bulletin Zoologisch Museum, Universiteit van Amsterdam* 11: 77-79.
- Verco, J.C. 1907. Notes on South Australian marine Mollusca, with descriptions of new species. Part VII. *Transactions of the Royal Society of South Australia* 31: 305-315.
- Verco, J.C. 1909. Notes on South Australian Mollusca, with descriptions of new species. Part X. *Transactions of the Royal Society of South Australia* 33: 270-276.
- Verco, J.C. 1916. Note on *Edentellina typica*, Gatliff and Gabriel. *Transactions of the Royal Society of South Australia* 40: 596-597.
- Verrill, A. E., & Emerton. 1882. Catalogue of marine Mollusca added to the fauna of the New England Region, during the past ten years. *Transactions of the Connecticut Academy of Science* 5(2): 447-587.
- Vonnemann, V., Schrödl, M., Klussmann-Kolb, A., & Wägele, H. 2005. Reconstruction of the phylogeny of the Opisthobranchia (Mollusca: Gastropoda) by means of 18S and 28S rRNA gene sequences. *Journal of Molluscan Studies* 71: 113-125.
- Wägele, H., & Klussman-Klob, A. 2005. Opisthobranchia (Mollusca, Gastropoda) - more than just slimy slugs. Shell reduction and its implications on defense and foraging. *Frontiers in Zoology* 2: 1-18.
- Wägele, H., Vonnemann, V., & Rudman, W. B. 2006. *Umbraculum umbraculum* (Lightfoot, 1786) (Gastropoda, Opisthobranchia, Tylodinoidea) and the synonymy of *U. mediterraneum* (Lamarck, 1812). *Records of the Western Australian Museum*, Supplement 69: 69-82.
- Wägele, H., & Willan, R. C. 2000. On the phylogeny of the Nudibranchia. *Zoological Journal of the Linnean Society* 130: 83-181.
- Watson, R. B. 1886. Mollusca of H.M.S. "Challenger" Expedition. Parts 18-20. *Journal of the Linnean Society of London, Zoology* 17(10): 284-293, 319-346.
- Wells, F. E. 1985. The taxonomic status of the opisthobranch mollusc *Bulla tenuissima* Sowerby, 1868. *Journal of the Malacological Society of Australia* 7(1-2): 29-33.
- Wells, F. E. 1989. Holoplanktonic molluscs (Class Gastropoda). Pp. 823-840 in: Shepherd, S.A. & Thomas, I.M. (eds), *Marine Invertebrates of Southern Australia. Part II*. South Australian Government Printing Division: Adelaide.
- Wells, F. E., & Bryce, C. W. 1993. *Sea Slugs of Western Australia*. Western Australian Museum: Perth.
- Willan, R. C. 1978. The nomenclature of three Pacific *Bulla* species. *Journal of the Malacological Society of Australia* 4: 57-68.
- Willan, R. C. 1983. New Zealand side-gilled sea slugs (Opisthobranchia: Notaspidea: Pleurobranchidae). *Malacologia* 23(2): 221-270.
- Willan, R. C. 1984. A review of diets in the Notaspidea (Mollusca: Opisthobranchia). *Journal of the Malacological Society of Australia* 6(3-4): 125-142.
- Willan, R. C. 1987a. Phylogenetic systematics of the Notaspidea (Opisthobranchia) with reappraisal of families and genera. *American Malacological Bulletin* 5(2): 215-241.
- Willan, R. C. 1987b. Phylogenetic systematics and zoogeography of Australian nudibranchs. 1. Presence of the aeolid *Godiva quadricolor* (Barnard) in Western Australia. *Journal of the Malacological Society of Australia* 8: 71-85.
- Willan, R. C. 1988. The taxonomy of two host-specific, cryptic dendronotoid nudibranch species (Mollusca: Gastropoda) from Australia including a new species description. *Zoological Journal of the Linnean Society* 94: 39-63.
- Willan, R. C. 1998. Orders Anaspidea and Notaspidea. Pp. 974-980 in: Beesley, P.L., Ross, G.J.B. & Wells, A. (eds), *Mollusca: The Southern Synthesis. Fauna of Australia. Volume 5*. CSIRO Publishing: Melbourne.
- Willan, R. C., & Bertsch, H. 1987. Description of a new pleurobranch (Opisthobranchia: Notaspidea) from Antarctic waters, with a review of notaspideans from southern polar seas. *Veliger* 29(3): 292-302.
- Willan, R. C., & Burn, R. 2003. On the publication date, authorship, and type species of *Umbraculum* and *Tylodina* (Gastropoda: Opisthobranchia: Tylodinoidea): a rejoinder. *Nautilus* 117: 23-29.
- Willan, R. C., & Coleman, N. 1984. *Nudibranchs of Australasia*. Australasian Marine Photographic Index: Caringbah, New South Wales. 56 pp.
- Wilson, J.B. 1887. List of Gastropoda, Lamellibranchiata, and Brachiopoda obtained at or near Port Phillip Heads. *Victorian Naturalist* 4: 116-118.
- Wilson, J.B. 1890. On some new species of marine molluscs. *Proceedings of the Royal Society of Victoria* 2: 64-67.
- Wilson, N.G. 2002. Egg masses of chromodorid nudibranchs (Mollusca: Gastropoda: Opisthobranchia). *Malacologia* 44: 289-305.
- Wilson, N.G. 2006. New record of the nudibranch *Polycera hedgpethi* Er. Marcus, 1964, in South Australia, with a discussion on its occurrence in South Australia. *Records of the Western Australian Museum*, Supplement 69: 137-140.
- Wilson, N.G., & Lee, M.S.Y. 2005. Molecular phylogeny of *Chromodoris* (Mollusca, Nudibranchia) and the identification of a planar spawning clade. *Molecular Phylogenetics and Evolution* 36: 722-727.